

Review

Immunomodulatory Role of Stem Cells in Cancer: A Landscape of Tumor-Promoting and Tumor-Inhibitory Effects

Manal Hadi Ghaffoori Kanaan ^{1,*}, Beom-Jin Lee ^{2,3}, Sura Saad Abdullah ², Chulhun Park ⁴, Abdolmajid Ghasemian ⁵, Marianne Marianne ⁶

1. Department of Food Industries/Technical Institute of Suwaria, Middle Technical University, Baghdad, Iraq; E-Mail: dr.manalhadi@mtu.edu.iq; ORCID: 0000-0002-9409-4778
2. College of Pharmacy, Ajou University, Suwon 16499, Republic of Korea; E-Mails: bjl@ajou.ac.kr; surasaad@ajou.ac.kr
3. Research Institute of Pharmaceutical Sciences and Technology, Ajou University, Suwon 16499, Republic of Korea
4. College of Pharmacy and Jeju Research Institute of Pharmaceutical Sciences, Jeju National University, Jeju 63243, Republic of Korea; E-Mail: chpark@jejunu.ac.kr
5. Noncommunicable Diseases Research Center, Fasa University of Medical Sciences, Fasa, Iran; E-Mail: majidghasemian86@gmail.com
6. Department of Pharmacology, Faculty of Pharmacy, Universitas Sumatera Utara, Medan, Indonesia; E-Mail: marianne80@usu.ac.id

* **Correspondence:** Manal Hadi Ghaffoori Kanaan; E-Mail: dr.manalhadi@mtu.edu.iq; ORCID: 0000-0002-9409-4778

Academic Editor: Lunawati L Bennett

OBM Genetics

2026, volume 10, issue 4

doi:10.21926/obm.genet.2604364

Received: July 17, 2026

Accepted: October 25, 2026

Published: October 30, 2026

Abstract

Stem cell therapy has emerged as a transformative frontier in cancer treatment, harnessing the unique immunomodulatory capabilities of stem cells to reshape anti-tumor immune responses. Mesenchymal stem cells (MSCs), hematopoietic stem cells (HSCs), induced pluripotent stem cells (iPSCs), and cancer stem cells (CSCs) dynamically interact with innate and adaptive immune systems within the tumor microenvironment (TME). MSCs secrete cytokines such as IL-6, TGF- β , and VEGF, which exhibit context-dependent effects on cytotoxic



© 2026 by the author. This is an open access article distributed under the conditions of the [Creative Commons by Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is correctly cited.

T lymphocytes (CTLs), regulatory T cells (Tregs), and myeloid-derived suppressor cells (MDSCs). These effects vary according to factors such as MSC tissue origin, tumor type, inflammatory status, and the cytokine milieu. MSCs can also modulate tumor-associated macrophage (TAM) polarization, but the direction of polarization toward pro-tumor M2-like or anti-tumor M1-like phenotypes is highly dependent on the surrounding TME. Conversely, CSCs drive immune evasion by expressing checkpoint molecules (e.g., PD-L1) and recruiting immunosuppressive cells, illustrating stem cells' paradoxical roles in cancer progression. Despite their promise, challenges persist: the TME's complexity, safety concerns (tumorigenicity, immune rejection), and a lack of standardized protocols impede clinical translation. Innovations such as genetically engineered stem cells for targeted drug delivery, their potential combination with immunotherapies such as CAR-T cells and checkpoint inhibitors, and personalized approaches tailored to tumor-specific profiles offer promising experimental strategies to enhance efficacy. This review underscores stem cells' potential role in rebalancing immune responses while emphasizing the need for further research to refine their therapeutic application. Although hematopoietic stem cell transplantation is clinically established for selected hematological malignancies, most other stem-cell-based immunomodulatory approaches remain preclinical or are under early clinical investigation.

Keywords

Stem cells; cancer; immunotherapy; immunomodulation; mesenchymal stem cells

1. Introduction

Stem cell therapy represents a frontier in regenerative medicine and cancer treatment, offering novel strategies to address the limitations of conventional therapeutic approaches [1]. Stem cells can self-renew and differentiate into diverse cell types, making them valuable for tissue repair, regenerative medicine, and targeted drug delivery [2]. Their ability to modulate the immune system has garnered considerable attention, particularly in cancer therapy [3].

Stem cells are undifferentiated cells characterized by self-renewal, enabling them to replicate and maintain a pool of undifferentiated cells, and differentiation, allowing them to mature into specialized cell types with specific functions. This dual capacity is fundamental to their role in tissue homeostasis, repair, and regeneration. Several types of stem cells with unique characteristics have therapeutic applications [4]. Importantly, mesenchymal stem cells (MSCs), hematopoietic stem cells (HSCs), induced pluripotent stem cells (iPSCs), and cancer stem cells (CSCs) have fundamentally different roles in cancer immunity: MSCs mainly exert paracrine immunomodulatory effects, HSCs generate hematopoietic and immune-cell lineages, iPSCs provide a renewable source for generating specialized immune cells, whereas CSCs are malignant stem-like cells involved in immune evasion and therapeutic resistance [4].

Stem cell therapy has broad applications, including regenerative medicine, where stem cells are employed to repair or replace damaged tissues and organs, offering hope for treating conditions such as heart disease, neurodegenerative disorders, and spinal cord injuries [5]. In cancer therapy, stem cells can serve as therapeutic carriers for targeted drug delivery to tumor sites, selectively

target CSCs to eradicate the root of the tumor, and modulate the immune system to enhance anti-tumor immune responses [2, 6]. Immunomodulation in cancer therapy involves modulating the immune system to better recognize and eliminate cancer cells, shifting the balance from tumor-promoting inflammation to anti-tumor immunity [7, 8]. This can be achieved by stimulating anti-tumor immune responses, suppressing immunosuppressive mechanisms that protect cancer cells from immune attack, and restoring immune homeostasis to create an environment less conducive to tumor growth [9]. Stem cells, particularly MSCs, play a pivotal role in immunomodulation by interacting with various immune cells, secreting cytokines and growth factors that influence immune cell behavior, and influencing immune cell differentiation and function, shaping the immune landscape within the tumor microenvironment (TME) [6, 10]. By modulating the immune system, stem cells can improve cancer therapy efficacy, overcome therapeutic resistance mechanisms used by cancer cells, and improve patient outcomes, offering a more personalized, targeted approach to cancer treatment [8].

1.1 Literature Search Strategy

This narrative review was based on literature searches conducted in PubMed, Scopus, and Google Scholar using combinations of the terms “stem cells,” “mesenchymal stem cells,” “hematopoietic stem cells,” “induced pluripotent stem cells,” “cancer stem cells,” “cancer,” “immunomodulation,” “tumor microenvironment,” “T cells,” “NK cells,” “macrophages,” “regulatory T cells,” “myeloid-derived suppressor cells,” and “immune evasion.” We considered studies published primarily between 2009 and 2026, and included relevant earlier studies when necessary. We included original studies, clinical studies, and relevant reviews on stem-cell-mediated cancer immunomodulation, and excluded studies without a clear link to cancer immunity or outside the scope of this review.

2. Cross-Talk of Stem Cells and the Immune System in Cancer

2.1 Overview of Immune Responses in Cancer

The immune system plays a complex and multifaceted role in cancer, characterized by both protective and promoting effects that significantly influence tumor development and progression [11]. Understanding the intricate interplay between innate and adaptive immunity is paramount for devising effective immunotherapeutic strategies that harness the immune system’s anti-tumor potential.

Innate immunity constitutes the body’s initial line of defense against pathogens and aberrant cells, including cancer cells, and involves a diverse array of cells such as macrophages, natural killer (NK) cells, and dendritic cells (DCs) [12]. Macrophages, versatile phagocytes, can engulf and digest cancer cells, while also presenting tumor-associated antigens to adaptive immune cells, thereby initiating a targeted anti-tumor response. NK cells, specialized cytotoxic lymphocytes, can directly recognize and kill cancer cells without prior sensitization, providing rapid and broad-spectrum anti-tumor activity [13]. DCs function as professional antigen-presenting cells (APCs), capturing, processing, and presenting tumor-associated antigens to T cells, thereby bridging the innate and adaptive immune responses and initiating a targeted anti-tumor response [14]. Myeloid-derived suppressor cells (MDSCs), a heterogeneous population of immature myeloid cells, represent

another component of the innate immune system, but they typically exert immunosuppressive effects within the TME, suppressing anti-tumor immune responses and promoting tumor growth and metastasis [15].

Adaptive immunity, a more refined and targeted defense mechanism, involves T and B lymphocytes, which recognize specific tumor-associated antigens and mount tailored immune responses to eradicate cancer cells [16]. T cells, including cytotoxic T lymphocytes (CTLs) and helper T cells, directly kill cancer cells and coordinate immune responses by secreting cytokines that modulate other immune cells [6]. B cells, responsible for humoral immunity, produce antibodies that can neutralize tumor antigens, activate the complement system to destroy cancer cells, and mediate antibody-dependent cell-mediated cytotoxicity (ADCC), recruiting immune cells to eliminate antibody-coated cancer cells. Regulatory T cells (Tregs), a subset of T cells, suppress immune responses to maintain immune homeostasis and prevent autoimmunity, but in cancer, they can also inhibit anti-tumor immunity within the TME, protecting cancer cells from immune attack [17].

2.2 Interaction Between Stem Cells and Immune Cells

Different stem cell populations interact with immune cells, influencing the balance between anti-tumor immunity and immune suppression and thereby affecting tumor growth and progression [18, 19]. For MSCs, these interactions are mainly mediated by cell-cell contact, cytokine secretion, and other signaling mechanisms and can either stimulate or suppress immune-cell activity depending on the surrounding microenvironment [20]. HSCs primarily contribute through hematopoietic and immune-cell generation, whereas iPSCs can serve as a source of differentiated immune effector cells. CSCs, in contrast, predominantly interact with immune cells to facilitate immune evasion. For example, mesenchymal stem cells (MSCs), known for their immunomodulatory properties, can interact with NK cells, suppressing their proliferation, cytokine secretion, and cytotoxicity, thereby potentially dampening the innate immune response against cancer [21]. However, activated NK cells can also lyse MSCs, highlighting a complex bidirectional interaction in which NK cells can eliminate MSCs that may pose a threat. MSCs can modulate DC differentiation, maturation, and function, influencing their ability to present tumor-associated antigens to T cells and initiate a targeted adaptive immune response [20, 22].

Tumor-associated macrophages (TAMs), a prominent component of the TME, interact extensively with CSCs to promote tumor progression, maintain CSC stemness, and enhance their resistance to therapy [23]. TAMs can secrete growth factors, cytokines, and other factors that enhance CSC self-renewal, proliferation, and drug resistance, while CSCs recruit macrophages into the TME, creating a self-reinforcing loop that promotes tumor growth and immune evasion. The interactions between TAMs and CSCs exemplify the complex interplay between stem cells and immune cells within the TME, highlighting the challenges in developing effective cancer immunotherapies [24].

3. Mechanisms of Immunomodulation by Stem Cells in Cancer

Stem cells, particularly MSCs, exert immunomodulatory effects in the TME through multifaceted mechanisms, including the secretion of cytokines (e.g., IL-6, IL-8, VEGF, TGF- β , IL-10) and chemokines (e.g., MCP-1, MIP-1), direct cell-cell interactions, and recruitment of immune cells such

as neutrophils, macrophages, T cells, Tregs, and MDSCs [25, 26]. These secreted factors modulate immune responses by influencing immune cell differentiation, activation, and function: pro-inflammatory cytokines can enhance anti-tumor immunity by recruiting cytotoxic T cells and NK cells, while immunosuppressive cytokines like TGF- β and IL-10 suppress T cell activity and promote tolerance. MSCs also alter macrophage polarization, shifting the balance between anti-tumor M1 (pro-inflammatory) and pro-tumor M2 (anti-inflammatory) phenotypes, and regulate T cell function [25, 27, 28]. Importantly, MSC-mediated regulation of Tregs, MDSCs, macrophages, and effector lymphocytes is context-dependent and can either enhance or suppress anti-tumor immunity according to MSC origin, activation status, tumor type, and the surrounding cytokine milieu. The complexity and context-dependence of these interactions, including their impact on the inflammatory landscape and immune cell dynamics, underscore the need for further research to optimize stem cell-based therapies for enhancing anti-tumor efficacy while mitigating immune suppression in the TME [29] (Table 1).

Table 1 Studies on the immunomodulatory role of stem cells in cancer.

Author(s)	Stem cell type	Cancer type	Main findings	Reference
Yu et al.	HCT116 (colon cancer stem cells)	Colon Cancer	TP5 reduces stemness in HCT116 cells, enhancing the cytotoxic effect of oxaliplatin by decreasing stemness-related markers and altering Wnt/ β -catenin signaling.	[30]
Casbon et al.	HSC	Breast Cancer	Tumor-derived G-CSF directs hematopoietic stem cells to produce immunosuppressive neutrophils, contributing to immune evasion.	[31]
Pont et al.	HSC	Hematological Malignancies	The study demonstrates the immunotherapeutic potential of donor T cells post-alloSCT and the necessity of analyzing antigen distribution to avoid adverse effects.	[32]
Lullo et al.	HSC	Multiple Myeloma	IMiDs reduce pro-tumor T cell subsets in MM during autologous transplantation, suggesting an anti-inflammatory role.	[33]
Sayed et al.	HSC	Various Murine Tumors	Activation of myelopoiesis and MDSC differentiation in tumor models lead to immunosuppression.	[34]

Lee et al.	HSC	Ovarian Cancer	CD8α+ dendritic cells from HSCs enhance antitumor immune activity against ovarian cancer.	[35]
Rossnagl et al.	Osteoblasts	Various cancer	EDA-FN from osteoblasts increases arginase in myeloid-derived cells, promoting cancer growth via immune modulation.	[36]
Witte et al.	MSC	Lung Cancer	MSCs trigger immunomodulation through phagocytosis by monocytic cells, suggesting a mechanism for their therapeutic effects despite their rapid disappearance.	[37]
Wellhausen et al.	HSC	Acute Myeloid Leukemia, B cell lymphoma, Acute T cell leukemia	Development of epitope edited CD45 CAR T cells enables universal therapy against blood cancers; protects HSCs from CAR T cell recognition and destruction.	[38]
Rueff et al.	HSC	Plasma Cell Myeloma	Rapid recovery of NK cells after autologous HSCT correlates with protection from relapse in patients.	[39]
Zhang et al.	MSC	Various	Tumor-conditioned MSCs display hematopoietic differentiation, with a diminished Ca ²⁺ influx facilitating the conversion to immune cells like macrophages.	[40]
Giles et al.	MSC	Metastatic Tumors	HSPCs contribute to immunosuppression within pre-metastatic niches, impacting tumor dissemination.	[41]
Wu et al.	Hematopoietic Stem and Progenitor Cells	Various cancers	The study reveals that HSPCs in cancer patients are myeloid-biased, contributing to the accumulation of MDSCs and disease progression.	[42]

3.1 Cytokine and Growth Factor Secretion

Stem cells exert significant immunomodulatory effects in cancer by secreting cytokines and growth factors, soluble mediators that influence immune cells, tumor cells, and other TME components, ultimately affecting tumor growth and the overall immune response [43]. These secreted factors can act locally within the TME or systemically, influencing immune cell trafficking, differentiation, and function.

Cytokines are signaling molecules that mediate communication between cells and play a crucial role in regulating immune responses, orchestrating the complex interactions among immune cell types and influencing the balance between pro-inflammatory and anti-inflammatory responses [44]. Stem cells, particularly MSCs, secrete a wide array of cytokines that can either promote or suppress immune responses, depending on the specific context and the target cells involved. For instance, MSCs secrete IL-6, which can have dual effects depending on the context, promoting tumor growth and angiogenesis in some cases, while contributing to inflammatory immune activation under other conditions [45, 46]. The net effect depends on tumor type, the cellular source of IL-6, its concentration, and activation of downstream pathways such as STAT3. Similarly, IL-8 secreted by MSCs can attract neutrophils and other immune cells to the TME, modulating the inflammatory landscape and potentially influencing tumor progression [47].

TGF- β is another important cytokine secreted by MSCs, known for its immunosuppressive properties, inhibiting the activation and proliferation of T cells, promoting the differentiation of Tregs, and suppressing the activity of NK cells. These effects can contribute to immune evasion by cancer cells. However, TGF- β can also have anti-tumor effects by inhibiting tumor-cell proliferation and inducing apoptosis, particularly during earlier stages of tumorigenesis, highlighting the complex and context-dependent nature of cytokine signaling in cancer [48].

Stem cells also secrete growth factors that influence tumor growth, angiogenesis, and immune cell function, shaping the TME and influencing the course of tumor development. VEGF is a key growth factor secreted by stem cells, promoting angiogenesis and tumor vascularization, providing tumors with the nutrients and oxygen they need to grow and metastasize. By promoting angiogenesis, VEGF can enhance tumor growth and metastasis. However, VEGF can also modulate immune cell function by suppressing the maturation of DCs and inhibiting T cell responses, further contributing to immune evasion [49, 50].

The specific effects of stem cell-derived cytokines on tumor growth and immune response depend on several factors, including stem cell type, the TME, and the specific cytokines involved, making it challenging to predict the overall impact of stem cell therapy on cancer progression [51]. Understanding these complex interactions is crucial for developing effective stem cell-based immunotherapies that can harness the beneficial effects of cytokines while minimizing their potential to promote tumor growth. For example, human Wharton's jelly stem cells (hWJSCs) secrete cytokines that inhibit the proliferation of ovarian cancer cells in vitro, suggesting their potential as a therapeutic agent. These hWJSC extracts decreased the expression of oncogenic cytokines, chemokines, and growth factors, thereby inhibiting OVCAR3 cells, demonstrating their ability to modulate the cytokine milieu within the TME [52].

3.2 Regulation of Tumor-Associated Macrophages

TAMs are a key component of the TME and play a critical role in tumor progression and immune evasion, influencing angiogenesis, metastasis, and therapeutic resistance [53]. Stem cells can influence macrophage polarization, affecting the balance between M1 and M2 macrophages, which have opposing roles in tumor immunity, with M1 macrophages generally exhibiting anti-tumor activity and M2 macrophages promoting tumor growth [54].

Macrophages exhibit remarkable plasticity, adapting their phenotype and function in response to environmental signals, and can differentiate into functionally distinct phenotypes, primarily classified as M1 and M2 macrophages [55]. M1 macrophages are classically activated by stimuli such as IFN- γ and lipopolysaccharide (LPS) and exert anti-tumor functions, producing pro-inflammatory cytokines such as IL-12 and TNF- α , mediating cytotoxicity, and promoting T cell responses. In contrast, M2 macrophages are alternatively activated by stimuli such as IL-4 and IL-10 and promote tumor progression, secreting anti-inflammatory cytokines such as IL-10 and TGF- β , promoting angiogenesis, and suppressing T cell responses [56].

MSCs can influence macrophage polarization in a context-dependent manner, potentially shifting macrophages toward either M1-like or M2-like phenotypes. For example, MSCs can secrete factors (e.g., TNF- α , IL-1 β , IL-6, IL-12, IL-23, and CXCL10) that promote M1 polarization, enhancing anti-tumor immunity and increasing pro-inflammatory cytokine production [57]. Conversely, stem cells can also promote M2 polarization, contributing to immune suppression and tumor progression by secreting factors that suppress T cell activity and promote angiogenesis [58]. Thus, the inflammatory environment, tumor-derived signals, and the functional state of MSCs determine MSC-mediated macrophage polarization.

TAMs play a critical role in tumor progression and immune evasion, influencing various aspects of tumor biology. M2 macrophages promote tumor growth by secreting growth factors and cytokines that stimulate tumor cell proliferation and angiogenesis, providing tumors with the resources they need to grow and metastasize [59]. They also suppress anti-tumor immune responses by inhibiting T cell activation and promoting the differentiation of Tregs, creating an immunosuppressive environment that protects cancer cells from immune attack. TAMs can also contribute to CSC maintenance and drug resistance, promoting tumor recurrence and therapeutic failure [60].

Targeting TAMs has emerged as a promising strategy for cancer immunotherapy, aiming to reprogram their function and enhance anti-tumor immunity. Strategies include depleting TAMs, inhibiting their recruitment to the TME, and reprogramming M2 macrophages to M1 macrophages, shifting the balance from tumor-promoting inflammation to anti-tumor immunity [61]. Preclinical approaches targeting innate immune signaling pathways, including STING activation, can also alter TAM function [24]. Stem cells can deliver therapeutic agents directly to TAMs, modulating their function and promoting anti-tumor immunity, offering a targeted approach to reprogram the TME. For instance, MSCs can deliver oncolytic Newcastle disease virus (NDV) to enhance therapeutic effects by altering the TME and promoting apoptosis, demonstrating stem cells' potential to modulate TAM function [62]. However, clinicians have not yet established durable MSC-mediated TAM reprogramming.

3.3 Modulation of T-Cell Responses

T cells are critical components of the adaptive immune system, recognizing and eliminating cancer cells through direct cytotoxicity and the secretion of cytokines that coordinate immune responses. Stem cells can influence T-cell activation, differentiation, and function, affecting their ability to mount effective anti-tumor responses and impacting the course of cancer progression [63].

T-cell activation is a complex process that requires multiple signals, including antigen presentation by APCs and co-stimulatory signals that provide the necessary cues for T-cell proliferation and effector function [64]. Stem cells can modulate T-cell activation by influencing APC function, altering the expression of co-stimulatory molecules (i.e., CD80/CD86, CD40, PD-L1, ICOS-L, and CTLA-4), and secreting cytokines that affect T-cell signaling pathways [65]. For example, MSCs can suppress T-cell proliferation and activation by secreting TGF- β and IL-10, inhibiting pro-inflammatory cytokine production and dampening the overall immune response [66].

Stem cells also influence T-cell differentiation, the process by which T cells acquire specialized functions, affecting the balance among T-cell subsets such as CTLs, helper T cells, and Tregs [67]. CTLs are critical for directly killing cancer cells, while helper T cells coordinate immune responses through cytokine secretion, activating other immune cells and enhancing anti-tumor immunity. Tregs, on the other hand, suppress immune responses to prevent autoimmunity but can also inhibit anti-tumor immunity in the TME, protecting cancer cells from immune attack [68].

Stem-cell-based approaches can modulate T-cell-mediated anti-tumor immunity, although the direction of this effect depends strongly on the stem cell type and experimental context. Engineered MSC-based strategies may promote CTL responses, whereas unmodified MSCs frequently suppress T-cell activation through immunoregulatory mediators. Similarly, their effects on Tregs may vary according to the cytokine environment and MSC activation state [69]. By modulating T-cell responses, stem-cell-based approaches may influence the efficacy of cancer therapies [70].

The potential therapeutic effects of stem cells on T-cell-mediated anti-tumor immunity have been demonstrated, providing evidence that they can enhance T-cell activity and promote tumor regression. For instance, MSCs loaded with oncolytic NDV can enhance E7-specific lymphocyte proliferation and CD8⁺ cytotoxicity, increasing the number of T cells targeting tumor-associated antigens and enhancing their cytotoxic activity [71]. Additionally, a strategy combining 4-methylumbelliferone (4Mu) with an adenovirus encoding interleukin-12 (AdIL-12) elicited a potent anti-tumor effect and prolonged animal survival by promoting cytotoxic-specific T cell responses, demonstrating the ability of stem cells to enhance T cell-mediated immunity [72].

3.4 Regulation of Regulatory T-Cells and Myeloid-Derived Suppressor Cells

As mentioned, Tregs and MDSCs are major players in immune suppression in the TME, inhibiting anti-tumor immune responses and promoting tumor growth and metastasis. These cells use various mechanisms to suppress immune cell activity, creating a protective shield around cancer cells and hindering immune elimination. Stem cells, particularly MSCs, can modulate the function of Tregs and MDSCs; however, these effects may be either suppressive or supportive depending on the surrounding inflammatory and tumor context [73]. Tregs can inhibit anti-tumor immunity, promoting tumor growth and immune evasion by suppressing the activity of effector T cells and other immune cells that would otherwise target and kill cancer cells [74]. Tregs suppress T-cell activation and proliferation via inhibitory cytokines (TGF- β , IL-10), metabolic disruption

(adenosine/CD39-CD73), and CTLA-4-mediated competition for co-stimulatory signals (CD80/CD86); inhibit NK cell activity by downregulating activating receptors (e.g., NKG2D) through TGF- β /IL-10 and inducing exhaustion via PD-1/PD-L1 or granzyme B; and promote MDSC differentiation through GM-CSF, CD40-CD40L interactions, and miRNA transfer, amplifying their arginase-1/iNOS-dependent immunosuppression. This establishes a self-reinforcing network within the TME, bolstered by crosstalk with M2-like macrophages, CAFs, and chemokine-driven recruitment of suppressive cells [75].

Also, MDSCs inhibit T cell activation and proliferation, promote Treg differentiation, and suppress NK cell activity, contributing to generalized immune suppression within the TME [76]. MDSCs also promote angiogenesis, tumor cell proliferation, and metastasis, facilitating tumor progression and dissemination to distant sites [77]. MSCs can regulate Tregs and MDSCs through several context-dependent mechanisms [78, 79]. Although some experimental conditions have been associated with reduced suppressive activity, MSC-derived factors such as TGF- β and IL-10 can also support Treg-mediated immunosuppression. Similarly, MSC effects on MDSCs vary according to tumor-derived signals and the inflammatory environment. Therefore, MSC-mediated regulation of Tregs and MDSCs should not be considered uniformly anti-tumor [79].

The mechanisms by which stem cells inhibit immune suppression in cancer are complex and context-dependent, requiring further investigation to fully elucidate their intricacies and optimize their therapeutic potential [80]. Understanding these mechanisms is crucial for designing effective stem cell-based immunotherapies that can selectively target Tregs and MDSCs while sparing other immune cells, maximizing anti-tumor immunity and minimizing off-target effects. For instance, oncolytic NDV delivered by MSCs can increase infiltration of CD11b⁺Gr1⁺ MDSCs, suggesting a modulation of the TME and a potential shift in MDSC function [81]. Additionally, SF2523, a dual PI3K/BRD4 inhibitor, blocks tumor immunosuppression and promotes adaptive immune responses by reducing the infiltration of MDSCs and the polarization of immunosuppressive macrophages, highlighting the potential of targeted therapies to modulate immune cell activity [82] (Figure 1).




Figure 1 Mechanisms of MSC-mediated immunomodulation in cancer. This schematic illustrates the context-dependent effects of mesenchymal stem cells (MSCs) on immune responses within the tumor microenvironment (TME). (a) Cytokine release: MSCs release interleukin-6 (IL-6), transforming growth factor-beta (TGF-β), vascular endothelial growth factor (VEGF), and interleukin-10 (IL-10). The predominant immunoregulatory effects of MSC-derived TGF-β include regulatory T-cell (Treg) differentiation and suppression of effector T-cell activity. IL-6 has context-dependent effects, whereas VEGF promotes angiogenesis and IL-10 contributes to immunosuppression. (b) Macrophage polarization: MSCs influence tumor-associated macrophage (TAM) polarization toward M1-associated anti-tumor activity or M2-associated angiogenesis and immunosuppression. (c) Context-dependent anti-tumor effects: Under specific conditions, MSC-mediated modulation may promote cytotoxic T-lymphocyte (CTL) activation and inhibit Treg activity; these effects are distinct from the predominant immunosuppressive effects of MSC-derived TGF-β. (d) Context-dependent MDSC modulation: MSCs may reduce myeloid-derived suppressor cell (MDSC)-mediated T-cell suppression under specific conditions. Solid arrows indicate activation or differentiation, inhibitory lines indicate suppression, and dashed arrows indicate indirect or context-dependent immunomodulatory effects. Context-dependent effects are additionally identified by explicit labels. Abbreviations: MSC, mesenchymal stem cell; TME, tumor microenvironment; IL, interleukin; TGF-β, transforming growth factor-beta; VEGF, vascular endothelial growth factor; TAM, tumor-associated macrophage; Treg, regulatory T cell; CTL, cytotoxic T lymphocyte; MDSC, myeloid-derived suppressor cell.

4. Types of Stem Cells Enrolled in Cancer Immunomodulation

4.1 Mesenchymal Stem Cells

MSCs are multipotent stromal cells that can differentiate into various cell types, including osteocytes, chondrocytes, and adipocytes, making them valuable for regenerative medicine and tissue engineering [83]. MSCs possess inherent immunomodulatory properties, making them attractive candidates for cell-based therapies in cancer, where they can modulate the immune system to enhance anti-tumor responses [84]. MSCs exert their immunomodulatory effects through diverse mechanisms, including the secretion of cytokines, direct cell-cell interactions with immune cells, and the modulation of immune cell function, creating a complex interplay between MSCs and the immune system [85]. MSCs can secrete immunosuppressive cytokines such as TGF- β and IL-10, which inhibit T cell proliferation and activation, dampening immune responses and promoting immune tolerance [86]. They can also modulate immune cells such as macrophages, NK cells, and DCs, influencing their activity and shaping the immune landscape within the TME [87].

Studies on MSC-based cancer immunotherapy have shown promising results predominantly in preclinical settings, demonstrating their potential to modulate the immune system and enhance anti-tumor responses. MSCs can be used to deliver therapeutic agents directly to tumor sites and modulate the immune system [83]. However, unmodified MSCs may also promote immunosuppression, angiogenesis, and tumor progression in certain tumor microenvironments; therefore, their anti-tumor versus pro-tumor effects depend on MSC origin, phenotype, tumor type, and cytokine exposure [88].

4.2 Hematopoietic Stem Cells

HSCs are multipotent stem cells that give rise to all blood cell types, including immune cells, making them essential for immune system function and reconstitution following damage or disease [89]. HSCs are primarily used in hematopoietic stem cell transplantation (HSCT) to restore the immune system in patients with hematological cancers, such as leukemia and lymphoma, where the patient's own immune system has been compromised by disease or treatment [90]. HSCT represents an established clinical application of stem cells in oncology, unlike many other stem-cell-based immunomodulatory strategies that remain experimental.

The immunotherapy potential of HSCs in hematological cancers lies in their ability to reconstitute the immune system and promote tumor rejection, allowing for the establishment of a new, healthy immune system that can target and eliminate cancer cells [91]. Allogeneic HSCT, where HSCs are derived from a donor, can induce a graft-versus-tumor (GVT) effect, where donor-derived immune cells recognize and kill tumor cells, providing a potent anti-tumor response [92]. HSCs can also be genetically modified to express T-cell receptors (TCRs) or chimeric antigen receptors (CARs) that target tumor-associated antigens, enhancing their ability to recognize and kill cancer cells [93].

4.3 Induced Pluripotent Stem Cells

Induced pluripotent stem cells (iPSCs) are generated from adult somatic cells through reprogramming, a process that reverses their differentiation and returns them to an embryonic stem cell-like state, capable of differentiating into any cell type in the body [94]. iPSCs offer

significant potential for personalized therapies, as they can be derived from a patient's own cells, reducing the risk of immune rejection and enabling patient-specific cell therapies [95].

Emerging research on iPSCs in immunotherapy focuses on their potential to generate immune cells with anti-tumor activity, creating a renewable source of immune cells that can be tailored to target specific cancers [96]. iPSCs can be differentiated into NK cells, T cells, and DCs that can be used to target and kill cancer cells, offering a versatile platform for developing novel immunotherapies [97]. Recent advances in NK-cell expansion, activation, and genetic engineering have further strengthened the potential of stem-cell-derived NK-cell platforms for cancer immunotherapy [98]. However, their clinical application remains under investigation. The mechanisms of immune modulation in cancer therapy using iPSCs are still being investigated, but their potential to generate personalized immune cell therapies holds great promise [96].

4.4 Cancer Stem Cells

CSCs are a subpopulation of tumor cells with stem-like properties that contribute to tumor initiation, progression, and therapeutic resistance, making them a critical target for cancer therapy [99]. CSCs can evade immune detection and elimination, contributing to tumor recurrence and metastasis, highlighting the need for strategies to overcome their immune evasion mechanisms [100].

The role of CSCs in immune evasion and therapeutic resistance has garnered significant attention, improving investigations into their interactions with the immune system and the mechanisms by which they escape immune surveillance [101]. CSCs can express high levels of immune checkpoint molecules, such as PD-L1, which inhibit T cell activation, preventing T cells from effectively targeting and killing cancer cells [102]. They can also secrete factors that suppress immune cell function and promote immune tolerance, creating an immunosuppressive environment that protects them from immune attack [103]. Additional mechanisms include reduced antigen presentation, expression of anti-phagocytic molecules such as CD47, and recruitment or activation of immunosuppressive populations including Tregs, MDSCs, and TAMs. These mechanisms can contribute to persistence of CSCs after treatment and thereby promote recurrence and therapeutic resistance [23].

Targeting CSCs has emerged as a promising strategy for cancer therapy, aiming to eliminate the root of the tumor and prevent recurrence and metastasis [6]. Strategies include developing drugs that specifically target CSCs, using immunotherapies to target CSC-associated antigens, blocking immune-evasion pathways such as PD-1/PD-L1 and CD47 signaling where appropriate, and modulating the TME to reduce CSC-mediated immune suppression. For instance, 4Mu decreases CD47 expression on hepatic cancer stem cells and primes a potent antitumor T cell response induced by IL-12, demonstrating the potential of targeting CSCs to enhance anti-tumor immunity (Figure 2) [104].




Figure 2 Cellular sources and pathways relevant to cancer immunomodulation. Somatic cells can be isolated and reprogrammed into induced pluripotent stem cells (iPSCs). Through established, protocol-dependent directed differentiation, iPSCs can generate immune cells, including T cells and macrophages, or differentiate into mesenchymal stem cells (MSCs). MSCs can also be isolated directly for potential therapeutic applications. Hematopoietic stem cells (HSCs) and hematopoietic stem/progenitor cells (HSPCs), obtained through bone marrow aspiration, give rise to immune cells through hematopoietic differentiation. MSCs regulate immune cells through immunomodulation and paracrine signaling rather than direct differentiation into T cells or macrophages. MSC-derived paracrine effects on tumor cells are context-dependent, while immune cells influence tumor cells through immune-mediated mechanisms. Solid arrows indicate activation or differentiation, inhibitory lines indicate suppression, and dashed arrows indicate indirect or context-dependent immunomodulatory effects. Solid arrows labeled “Isolation,” “Bone marrow aspiration,” or “Reprogramming” denote the corresponding procedures; the MSC-to-tumor-cell arrow represents the explicitly labeled context-dependent paracrine effects. Abbreviations: iPSC, induced pluripotent stem cell; MSC, mesenchymal stem cell; HSC, hematopoietic stem cell; HSPC, hematopoietic stem/progenitor cell.

5. Challenges and Limitations

5.1 Tumor Microenvironment and Immune Evasion

The TME critically influences the effectiveness of stem cell-based immunomodulation [105]. The TME is a complex network of cells, including immune cells, fibroblasts, and endothelial cells, along with extracellular matrix components and soluble factors. The TME can promote tumor growth and metastasis by suppressing anti-tumor immune responses and promoting angiogenesis [106]. Immune evasion is a major challenge in cancer therapy. Cancer cells can evade immune detection

and elimination through various mechanisms, including the expression of immune checkpoint molecules, the secretion of immunosuppressive factors, and the recruitment of immunosuppressive cells [107]. The TME can also contribute to immune evasion by creating a physical barrier that prevents stem cells and stem cell-derived immune cells from reaching tumor cells [108].

5.2 Safety and Ethical Concerns

The safety of stem cell-based immunotherapies is a major concern [109]. Potential risks include tumorigenicity or teratoma formation from residual undifferentiated pluripotent cells, unwanted differentiation, immune rejection, transmission of infectious agents, genomic instability after prolonged culture or genetic manipulation, and potential enhancement of tumor growth by some MSC populations. Immune rejection can occur with allogeneic cells, while infectious-agent transmission remains a risk if cells are not properly screened and handled [110]. Long-term safety also remains incompletely defined for many genetically modified or stem-cell-derived therapeutic products.

Ethical issues surrounding stem cell use in cancer therapy include informed consent, patient selection, and the potential for commercial exploitation. Patients must be fully informed about the potential risks and benefits of stem cell therapy before undergoing treatment. Patient selection criteria must be carefully considered to ensure that patients are likely to benefit from stem cell therapy. The commercialization of stem cell therapy must be carefully regulated to prevent exploitation of vulnerable patients [111].

Recent preclinical and clinical development has attempted to reduce these risks through stringent cell characterization, elimination of residual undifferentiated cells, genomic-stability testing, standardized differentiation and purification procedures, sterility testing, controlled dosing, and long-term follow-up. Gene-engineered cell products may also incorporate safety switches to permit elimination of the administered cells if severe toxicity develops [112].

5.3 Regulatory Issues and Standardization

Regulatory challenges in stem cell-based cancer therapies include unclear regulatory guidelines and the need for standardized protocols. Stem cell therapies are often classified as biologics, which require rigorous testing and approval processes. However, the regulatory pathways for stem cell therapies are not always clear, leading to delays in the development and approval of new therapies [113].

Standardization of protocols for stem cell isolation and therapy is essential to ensure the reproducibility and reliability of stem-cell-based approaches. Standardized protocols are needed for stem-cell isolation, culture, characterization, and delivery. Use predefined quality-control criteria to assess cell identity, purity, viability, potency, genomic stability, sterility, and batch-to-batch consistency. Manufacturing processes should also control donor or starting-material selection, culture conditions, passage number, cryopreservation, and, when applicable, genetic modification. These requirements are particularly important for clinical-grade products manufactured under Good Manufacturing Practice conditions [114].

6. Future Directions and Perspectives

Genetic modification of stem cells can enhance their immunomodulatory properties and improve therapeutic efficacy. Stem cells can be engineered to express anti-tumor cytokines, deliver oncolytic viruses, or target cancer stem cells [115]. For example, CAR T-cells can be engineered for adoptive cell therapy of cancer using long-term episomal gene transfer [116].

Combining stem-cell-based approaches with immune checkpoint inhibitors or CAR-based immune-cell therapies has been proposed as a strategy to improve anti-tumor activity. Potential mechanisms include modulating the immunosuppressive TME, improving delivery of therapeutic molecules, and generating stem-cell-derived immune effector cells. However, most evidence for these combinations remains preclinical, and major barriers include limited cell persistence and tumor infiltration, tumor heterogeneity, immunosuppressive cytokines, manufacturing complexity, and safety concerns [117, 118].

Recent developments have also expanded the field toward engineered NK-cell therapies and advanced stem-cell-derived experimental models. NK-cell expansion, activation, and genetic engineering are being investigated to improve persistence and tumor cytotoxicity [119], while stem-cell-derived organoid systems provide increasingly sophisticated models for studying tissue development, disease, and therapeutic responses [98, 120].

Tailoring stem-cell-based therapies to individual patient needs and tumor characteristics may improve therapeutic efficacy and reduce side effects. Personalized approaches may involve selecting the most appropriate cellular platform, genetically modifying cells to target specific tumor antigens, and optimizing their delivery to tumor sites [121]. However, such personalized strategies remain largely investigational and require validation in prospective clinical studies.

7. Conclusion

Stem cells and stem-cell-derived products can exert diverse immunomodulatory effects in cancer, influencing immune cells and the TME. These effects include cytokine secretion, modulation of macrophage polarization, and regulation of T-cell responses, particularly in the case of MSCs. However, these effects are highly context-dependent and may either suppress or promote tumor progression. Hematopoietic stem cell transplantation represents an established clinical application for selected hematological malignancies, whereas most MSC-, iPSC-, and engineered stem-cell-based immunotherapeutic approaches remain experimental or in early clinical development. Major challenges include tumorigenicity, unwanted differentiation, genomic instability, immune rejection, manufacturing variability, and uncertain long-term safety. Future studies should therefore focus on clarifying mechanisms, standardizing manufacturing, rigorously evaluating clinical benefit, and identifying patients most likely to benefit from these approaches.

Acknowledgments

All authors declare there is no acknowledgment in this study.

Author Contributions

Manal Hadi Ghaffoori Kanaan: conceptualization, supervision, writing – original draft, writing – review & editing; Beom-Jin Lee: conceptualization, investigation, writing – original draft, writing –

review & editing; Sura Saad Abdullah: writing – original draft, investigation; Chulhun Park: writing – review & editing; Abdolmajid Ghasemian: writing – review & editing; Marianne Marianne: writing – review & editing. All the authors critically revised and approved the final version of the manuscript.

Funding

There is no financial support for this study.

Competing Interests

The authors declare no competing interests related to this research.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

AI-Assisted Technologies Statement

Artificial intelligence (AI) tools were used solely for basic grammar correction and language refinement in the preparation of this manuscript. Specifically, ChatGPT6.1 was employed to improve the readability and linguistic clarity of the English text. All scientific content, data interpretation, and conclusions were developed independently by the author. The authors have thoroughly reviewed and edited the AI-assisted text to ensure its accuracy and accept full responsibility for the content of the manuscript.

References

1. Zakrzewski W, Dobrzyński M, Szymonowicz M, Rybak Z. Stem cells: Past, present, and future. *Stem Cell Res Ther.* 2019; 10: 68.
2. Alatyat SM, Alasmari HM, Aleid OA, Abdel-Maksoud MS, Elsherbiny N. Umbilical cord stem cells: Background, processing and applications. *Tissue Cell.* 2020; 65: 101351.
3. Yamanaka S. Pluripotent stem cell-based cell therapy-promise and challenges. *Cell Stem Cell.* 2020; 27: 523-531.
4. Mazini L, Rochette L, Admou B, Amal S, Malka G. Hopes and limits of adipose-derived stem cells (ADSCs) and mesenchymal stem cells (MSCs) in wound healing. *Int J Mol Sci.* 2020; 21: 1306.
5. Lee SJ, Jung C, Oh JE, Kim S, Lee S, Lee JY, et al. Generation of red blood cells from human pluripotent stem cells-An update. *Cells.* 2023; 12: 1554.
6. Biserova K, Jakovlevs A, Uljanovs R, Strumfa I. Cancer stem cells: Significance in origin, pathogenesis and treatment of glioblastoma. *Cells.* 2021; 10: 621.
7. Yilmaz M, Kaplan F, Mender I, Gryaznov SM, Dikmen ZG. Cancer stem cells and anti-tumor immunity. *Curr Stem Cell Res Ther.* 2023; 18: 445-459.
8. Kode JA, Mukherjee S, Joglekar MV, Hardikar AA. Mesenchymal stem cells: Immunobiology and role in immunomodulation and tissue regeneration. *Cytotherapy.* 2009; 11: 377-391.
9. Al-Ghadban S, Bunnell BA. Adipose tissue-derived stem cells: Immunomodulatory effects and therapeutic potential. *Physiology.* 2020; 35: 125-133.

10. Najar M, Raicevic G, Fayyad-Kazan H, Bron D, Toungouz M, Lagneaux L. Mesenchymal stromal cells and immunomodulation: A gathering of regulatory immune cells. *Cytotherapy*. 2016; 18: 160-171.
11. Chuang YM, Tzeng SF, Ho PC, Tsai CH. Immunosurveillance encounters cancer metabolism. *EMBO Rep*. 2024; 25: 471-488.
12. Kunimasa K, Goto T. Immunosurveillance and immunoediting of lung cancer: Current perspectives and challenges. *Int J Mol Sci*. 2020; 21: 597.
13. Gopalakrishnan V, Helmink BA, Spencer CN, Reuben A, Wargo JA. The influence of the gut microbiome on cancer, immunity, and cancer immunotherapy. *Cancer Cell*. 2018; 33: 570-580.
14. Wang C, Barnoud C, Cenerenti M, Sun M, Caffa I, Kizil B, et al. Dendritic cells direct circadian anti-tumour immune responses. *Nature*. 2023; 614: 136-143.
15. Ma P, Beatty PL, McKolanis J, Brand R, Schoen RE, Finn OJ. Circulating myeloid derived suppressor cells (MDSC) that accumulate in premalignancy share phenotypic and functional characteristics with MDSC in cancer. *Front Immunol*. 2019; 10: 1401.
16. Rezaie J, Chodari L, Mohammadpour-Asl S, Jafari A, Niknam Z. Cell-mediated barriers in cancer immunosurveillance. *Life Sci*. 2024; 342: 122528.
17. Smyth MJ, Dunn GP, Schreiber RD. Cancer immunosurveillance and immunoediting: The roles of immunity in suppressing tumor development and shaping tumor immunogenicity. *Adv Immunol*. 2006; 90: 1-50.
18. Liu N, Xiao X, Zhang Z, Mao C, Wan M, Shen J. Advances in cancer vaccine research. *ACS Biomater Sci Eng*. 2023; 9: 5999-6023.
19. Wang F, Cheng F, Zheng F. Stem cell like memory T cells: A new paradigm in cancer immunotherapy. *Clin Immunol*. 2022; 241: 109078.
20. Wu R, Fan X, Wang Y, Shen M, Zheng Y, Zhao S, et al. Mesenchymal stem cell-derived extracellular vesicles in liver immunity and therapy. *Front Immunol*. 2022; 13: 833878.
21. Yuan H, Li S, Zhao Z, Wang Y. Regulation of interferon- β -modified human umbilical cord mesenchymal stem cell-derived exosomes in proliferation and apoptosis of prostate cancer cells. *Organogenesis*. 2023; 19: 2285836.
22. Dehnavi S, Dadmanesh M, Rouzbahani NH, Karimi M, Asadirad A, Gholami M, et al. Mesenchymal stem cell-derived exosome: An interesting nanocarrier to improve allergen-specific intranasal immunotherapy. *Iran J Allergy Asthma Immunol*. 2023; 22: 561-574.
23. Pu Y, Ji Q. Tumor-associated macrophages regulate PD-1/PD-L1 immunosuppression. *Front Immunol*. 2022; 13: 874589.
24. Miao L, Qi J, Zhao QI, Wu QN, Wei DL, Wei XL, et al. Targeting the STING pathway in tumor-associated macrophages regulates innate immune sensing of gastric cancer cells. *Theranostics*. 2020; 10: 498-515.
25. Elliott SL, Kose E, Lewis AL, Steinfeld AE, Zollinger EA. Modeling the stem cell hypothesis: Investigating the effects of cancer stem cells and TGF- β on tumor growth. *Math Biosci Eng*. 2019; 16: 7177-7194.
26. Nazari F, Pearson AT, Nör JE, Jackson TL. A mathematical model for IL-6-mediated, stem cell driven tumor growth and targeted treatment. *PLoS Comput Biol*. 2018; 14: e1005920.
27. Chanmee T, Ontong P, Mochizuki N, Kongtawelert P, Konno K, Itano N. Excessive hyaluronan production promotes acquisition of cancer stem cell signatures through the coordinated

- regulation of Twist and the transforming growth factor β (TGF- β)-Snail signaling axis. *J Biol Chem.* 2014; 289: 26038-26056.
28. Arabpour M, Saghaizadeh A, Rezaei N. Anti-inflammatory and M2 macrophage polarization-promoting effect of mesenchymal stem cell-derived exosomes. *Int Immunopharmacol.* 2021; 97: 107823.
29. Yang L, Shi P, Zhao G, Xu J, Peng W, Zhang J, et al. Targeting cancer stem cell pathways for cancer therapy. *Signal Transduct Target Ther.* 2020; 5: 8.
30. Yu PC, Liu D, Han ZX, Liang F, Hao CY, Lei YT, et al. Thymopentin-mediated inhibition of cancer stem cell stemness enhances the cytotoxic effect of oxaliplatin on colon cancer cells. *Front Pharmacol.* 2022; 13: 779715.
31. Casbon AJ, Reynaud D, Park C, Khuc E, Gan DD, Schepers K, et al. Invasive breast cancer reprograms early myeloid differentiation in the bone marrow to generate immunosuppressive neutrophils. *Proc Natl Acad Sci.* 2015; 112: E566-E575.
32. Pont MJ, Honders MW, Kremer AN, van Kooten C, Out C, Hiemstra PS, et al. Microarray gene expression analysis to evaluate cell type specific expression of targets relevant for immunotherapy of hematological malignancies. *PLoS One.* 2016; 11: e0155165.
33. Di Lullo G, Marcatti M, Heltai S, Tresoldi C, Paganoni AM, Bordignon C, et al. Immunomodulatory drugs in the context of autologous hematopoietic stem cell transplantation associate with reduced pro-tumor T cell subsets in multiple myeloma. *Front Immunol.* 2019; 9: 3171.
34. Al Sayed MF, Amrein MA, Bühner ED, Huguenin AL, Radpour R, Riether C, et al. T-cell-secreted TNF α induces emergency myelopoiesis and myeloid-derived suppressor cell differentiation in cancer. *Cancer Res.* 2019; 79: 346-359.
35. Lee SW, Lee H, Lee KW, Kim MJ, Kang SW, Lee YJ, et al. CD8 α ⁺ dendritic cells potentiate antitumor and immune activities against murine ovarian cancers. *Sci Rep.* 2023; 13: 98.
36. Rosnagl S, Altrock E, Sens C, Kraft S, Rau K, Milsom MD, et al. EDA-fibronectin originating from osteoblasts inhibits the immune response against cancer. *PLoS Biol.* 2016; 14: e1002562.
37. de Witte SF, Luk F, Sierra Parraga JM, Garghesha M, Merino A, Korevaar SS, et al. Immunomodulation by therapeutic mesenchymal stromal cells (MSC) is triggered through phagocytosis of MSC by monocytic cells. *Stem Cells.* 2018; 36: 602-615.
38. Wellhausen N, O'Connell RP, Lesch S, Engel NW, Rennels AK, Gonzales D, et al. Epitope base editing CD45 in hematopoietic cells enables universal blood cancer immune therapy. *Sci Transl Med.* 2023; 15: eadi1145.
39. Rueff J, Medinger M, Heim D, Passweg J, Stern M. Lymphocyte subset recovery and outcome after autologous hematopoietic stem cell transplantation for plasma cell myeloma. *Biol Blood Marrow Transplant.* 2014; 20: 896-899.
40. Zhang L, Tang A, Zhou Y, Tang J, Luo Z, Jiang C, et al. Tumor-conditioned mesenchymal stem cells display hematopoietic differentiation and diminished influx of Ca²⁺. *Stem Cells Dev.* 2012; 21: 1418-1428.
41. Giles AJ, Reid CM, Evans JD, Murgai M, Vicioso Y, Highfill SL, et al. Activation of hematopoietic stem/progenitor cells promotes immunosuppression within the pre-metastatic niche. *Cancer Res.* 2016; 76: 1335-1347.
42. Wu WC, Sun HW, Chen HT, Liang J, Yu XJ, Wu C, et al. Circulating hematopoietic stem and progenitor cells are myeloid-biased in cancer patients. *Proc Natl Acad Sci.* 2014; 111: 4221-4226.

43. Markopoulos GS, Roupakia E, Marcu KB, Kolettas E. Epigenetic regulation of inflammatory cytokine-induced epithelial-to-mesenchymal cell transition and cancer stem cell generation. *Cells*. 2019; 8: 1143.
44. Kuo MH, Chen PY, Yang YP, Zheng MY, Miao CC, Wen KC, et al. Cytokine and epigenetic regulation of programmed death-ligand 1 in stem cell differentiation and cancer cell plasticity. *Stem Cells*. 2021; 39: 1298-1309.
45. Radharani NN, Yadav AS, Nimma R, Kumar TS, Bulbule A, Chanukuppa V, et al. Tumor-associated macrophage derived IL-6 enriches cancer stem cell population and promotes breast tumor progression via Stat-3 pathway. *Cancer Cell Int*. 2022; 22: 122.
46. Ibrahim SA, Gadalla R, El-Ghonaimy EA, Samir O, Mohamed HT, Hassan H, et al. Syndecan-1 is a novel molecular marker for triple negative inflammatory breast cancer and modulates the cancer stem cell phenotype via the IL-6/STAT3, Notch and EGFR signaling pathways. *Mol Cancer*. 2017; 16: 57.
47. Wang J, Wang Y, Wang S, Cai J, Shi J, Sui X, et al. Bone marrow-derived mesenchymal stem cell-secreted IL-8 promotes the angiogenesis and growth of colorectal cancer. *Oncotarget*. 2015; 6: 42825-42837.
48. Patel SA, Meyer JR, Greco SJ, Corcoran KE, Bryan M, Rameshwar P. Mesenchymal stem cells protect breast cancer cells through regulatory T cells: Role of mesenchymal stem cell-derived TGF- β . *J Immunol*. 2010; 184: 5885-5894.
49. Pakravan K, Babashah S, Sadeghizadeh M, Mowla SJ, Mossahebi-Mohammadi M, Ataei F, et al. MicroRNA-100 shuttled by mesenchymal stem cell-derived exosomes suppresses in vitro angiogenesis through modulating the mTOR/HIF-1 α /VEGF signaling axis in breast cancer cells. *Cell Oncol*. 2017; 40: 457-470.
50. Wu M, Chen L, Qi Y, Ci H, Mou S, Yang J, et al. Human umbilical cord mesenchymal stem cell promotes angiogenesis via integrin β 1/ERK1/2/HIF-1 α /VEGF-A signaling pathway for off-the-shelf breast tissue engineering. *Stem Cell Res Ther*. 2022; 13: 99.
51. Yazdani Y, Rad MR, Taghipour M, Chenari N, Ghaderi A, Razmkhah M. Genistein suppression of matrix metalloproteinase 2 (MMP-2) and vascular endothelial growth factor (VEGF) expression in mesenchymal stem cell like cells isolated from high and low grade gliomas. *Asian Pac J Cancer Prev*. 2016; 17: 5303-5307.
52. Kalamegam G, Sait KH, Anfinan N, Kadam R, Ahmed F, Rasool M, et al. Cytokines secreted by human Wharton's jelly stem cells inhibit the proliferation of ovarian cancer (OVCAR3) cells in vitro. *Oncol Lett*. 2019; 17: 4521-4531.
53. Osman A, Oze M, Afify SM, Hassan G, El-Ghlban S, Nawara HM, et al. Tumor-associated macrophages derived from cancer stem cells. *Acta Histochem*. 2020; 122: 151628.
54. Li D, Zhang Q, Li L, Chen K, Yang J, Dixit D, et al. β 2-microglobulin maintains glioblastoma stem cells and induces M2-like polarization of tumor-associated macrophages. *Cancer Res*. 2022; 82: 3321-3334.
55. Luo S, Yang G, Ye P, Cao N, Chi X, Yang WH, et al. Macrophages are a double-edged sword: Molecular crosstalk between tumor-associated macrophages and cancer stem cells. *Biomolecules*. 2022; 12: 850.
56. Jafarzadeh N, Safari Z, Pornour M, Amirizadeh N, Forouzandeh Moghadam M, Sadeghizadeh M. Alteration of cellular and immune-related properties of bone marrow mesenchymal stem cells

- and macrophages by K562 chronic myeloid leukemia cell derived exosomes. *J Cell Physiol.* 2019; 234: 3697-3710.
57. Wolfe AR, Trenton NJ, Debeb BG, Larson R, Ruffell B, Chu K, et al. Mesenchymal stem cells and macrophages interact through IL-6 to promote inflammatory breast cancer in pre-clinical models. *Oncotarget.* 2016; 7: 82482-82492.
 58. Sun W, Yan S, Yang C, Yang J, Wang H, Li C, et al. Mesenchymal stem cells-derived exosomes ameliorate lupus by inducing M2 macrophage polarization and regulatory T cell expansion in MRL/lpr mice. *Immunol Invest.* 2022; 51: 1785-1803.
 59. Liu X, Zhang M, Liu H, Zhu R, He H, Zhou Y, et al. Bone marrow mesenchymal stem cell-derived exosomes attenuate cerebral ischemia-reperfusion injury-induced neuroinflammation and pyroptosis by modulating microglia M1/M2 phenotypes. *Exp Neurol.* 2021; 341: 113700.
 60. Weng YS, Tseng HY, Chen YA, Shen PC, Al Haq AT, Chen LM, et al. MCT-1/miR-34a/IL-6/IL-6R signaling axis promotes EMT progression, cancer stemness and M2 macrophage polarization in triple-negative breast cancer. *Mol Cancer.* 2019; 18: 42.
 61. Zhang Q, Tsui YM, Zhang VX, Lu AJ, Lee JM, Lee E, et al. Reciprocal interactions between malignant cells and macrophages enhance cancer stemness and M2 polarization in HBV-associated hepatocellular carcinoma. *Theranostics.* 2024; 14: 892-910.
 62. Keshavarz M, Ebrahimzadeh MS, Miri SM, Dianat-Moghadam H, Ghorbanhosseini SS, Mohebbi SR, et al. Oncolytic Newcastle disease virus delivered by mesenchymal stem cells-engineered system enhances the therapeutic effects altering tumor microenvironment. *Virol J.* 2020; 17: 64.
 63. Gebhardt T, Park SL, Parish IA. Stem-like exhausted and memory CD8⁺ T cells in cancer. *Nat Rev Cancer.* 2023; 23: 780-798.
 64. Ascic E, Pereira CF. Transcription factor-mediated reprogramming to antigen-presenting cells. *Curr Opin Genet Dev.* 2025; 90: 102300.
 65. Zhang R, Liu TY, Senju S, Haruta M, Hirose N, Suzuki M, et al. Generation of mouse pluripotent stem cell-derived proliferating myeloid cells as an unlimited source of functional antigen-presenting cells. *Cancer Immunol Res.* 2015; 3: 668-677.
 66. Shephard MT, Merkhani MM, Forsyth NR. Human mesenchymal stem cell secretome driven T cell immunomodulation is IL-10 dependent. *Int J Mol Sci.* 2022; 23: 13596.
 67. Wang D, Huang S, Yuan X, Liang J, Xu R, Yao G, et al. The regulation of the Treg/Th17 balance by mesenchymal stem cells in human systemic lupus erythematosus. *Cell Mol Immunol.* 2017; 14: 423-431.
 68. Nozari P, Mokhtari P, Nemati M, Zainodini N, Taghipour Z, Asadi F, et al. Investigation of the effect of IFN- γ /TNF- α -treated mesenchymal stem cells on Th9-and Treg cell-related parameters in a mouse model of ovalbumin-induced allergic asthma. *Immunopharmacol Immunotoxicol.* 2022; 44: 773-785.
 69. Luo Y, Guo J, Zhang P, Cheuk YC, Jiang Y, Wang J, et al. Mesenchymal stem cell protects injured renal tubular epithelial cells by regulating mTOR-mediated Th17/Treg axis. *Front Immunol.* 2021; 12: 684197.
 70. Court AC, Le-Gatt A, Luz-Crawford P, Parra E, Aliaga-Tobar V, B  t  z LF, et al. Mitochondrial transfer from MSCs to T cells induces Treg differentiation and restricts inflammatory response. *EMBO Rep.* 2020; 21: EMBR201948052.

71. Ghorbani Alvanegh A, Mirzaei Nodooshan M, Dorostkar R, Ranjbar R, Jalali Kondori B, Shahriary A, et al. Antiproliferative effects of mesenchymal stem cells carrying *Newcastle disease virus* and *Lactobacillus Casei* extract on CT26 Cell line: Synergistic effects in cancer therapy. *Infect Agents Cancer*. 2023; 18: 46.
72. Zhao WH, Cheng JX, Shi PF, Huang JY. Human umbilical cord mesenchymal stem cells with adenovirus-mediated interleukin 12 gene transduction inhibits the growth of ovarian carcinoma cells both *in vitro* and *in vivo*. *J South Med Univ*. 2011; 31: 903-907.
73. Rui K, Hong Y, Zhu Q, Shi X, Xiao F, Fu H, et al. Olfactory ecto-mesenchymal stem cell-derived exosomes ameliorate murine Sjögren's syndrome by modulating the function of myeloid-derived suppressor cells. *Cell Mol Immunol*. 2021; 18: 440-451.
74. Juin SK, Ghosh S, Majumdar S. Glycyrrhizic acid facilitates anti-tumor immunity by attenuating Tregs and MDSCs: An immunotherapeutic approach. *Int Immunopharmacol*. 2020; 88: 106932.
75. Rui X, Alvarez Calderon F, Wobma H, Gerdemann U, Albanese A, Cagnin L, et al. Human OX40L–CAR-T_{regs} target activated antigen-presenting cells and control T cell alloreactivity. *Sci Transl Med*. 2024; 16: eadj9331.
76. Liu J, Liu J, Qin G, Li J, Fu Z, Li J, et al. MDSCs-derived GPR84 induces CD8⁺ T-cell senescence via p53 activation to suppress the antitumor response. *J Immunother Cancer*. 2023; 11: e007802.
77. Zhao Z, Qin J, Qian Y, Huang C, Liu X, Wang N, et al. FFAR2 expressing myeloid-derived suppressor cells drive cancer immunoevasion. *J Hematol Oncol*. 2024; 17: 9.
78. Pettinella F, Mariotti B, Lattanzi C, Bruderek K, Donini M, Costa S, et al. Surface CD52, CD84, and PTGER2 mark mature PMN-MDSCs from cancer patients and G-CSF-treated donors. *Cell Rep Med*. 2024; 5: 101380.
79. Abo-Aziza FA, Wasfy BM, Wahba SM, Abd-Elhalem SS. Mesenchymal stem cells and myeloid-derived suppressor cells interplay in adjuvant-induced arthritis rat model. *Int Immunopharmacol*. 2023; 120: 110300.
80. He JY, Huo FY, Tang HC, Liu B, Bu LL. Myeloid-derived suppressor cells in head and neck squamous cell carcinoma. *Int Rev Cell Mol Biol*. 2023; 375: 33-92.
81. Keshavarz M, Ebrahimzadeh MS, Miri SM, Dianat-Moghadam H, Ghorbanhosseini SS, Mohebbi SR, et al. Correction to: Oncolytic Newcastle disease virus delivered by mesenchymal stem cells-engineered system enhances the therapeutic effects altering tumor microenvironment. *Virology*. 2020; 17: 175.
82. Joshi S, Singh AR, Liu KX, Pham TV, Zulcic M, Skola D, et al. SF2523: Dual PI3K/BRD4 inhibitor blocks tumor immunosuppression and promotes adaptive immune responses in cancer. *Mol Cancer Ther*. 2019; 18: 1036-1044.
83. Lan T, Luo M, Wei X. Mesenchymal stem/stromal cells in cancer therapy. *J Hematol Oncol*. 2021; 14: 195.
84. Tu Z, Karnoub AE. Mesenchymal stem/stromal cells in breast cancer development and management. *Semin Cancer Biol*. 2022; 86: 81-92.
85. Weng Z, Zhang B, Wu C, Yu F, Han B, Li B, et al. Therapeutic roles of mesenchymal stem cell-derived extracellular vesicles in cancer. *J Hematol Oncol*. 2021; 14: 136.
86. Fan M, Zhang Y, Shi H, Xiang L, Yao H, Lin R. Bone mesenchymal stem cells promote gastric cancer progression through TGF- β 1/Smad2 positive feedback loop. *Life Sci*. 2023; 323: 121657.

87. Zhang H, Cao X, Gui R, Li Y, Zhao X, Mei J, et al. Mesenchymal stem/stromal cells in solid tumor microenvironment: Orchestrating NK cell remodeling and therapeutic insights. *Int Immunopharmacol.* 2024; 142: 113181.
88. Moreno R. Mesenchymal stem cells and oncolytic viruses: Joining forces against cancer. *J Immunother Cancer.* 2021; 9: e001684.
89. Gratwohl A, Baldomero H, Demirer TO, Rosti G, Dini G, Ladenstein R, et al. Hematopoietic stem cell transplantation for solid tumors in Europe. *Ann Oncol.* 2004; 15: 653-660.
90. Madjd Z, Gheytaichi E, Erfani E, Asadi-Lari M. Application of stem cells in targeted therapy of breast cancer: A systematic review. *Asian Pac J Cancer Prev.* 2013; 14: 2789-2800.
91. Gunduz BO, Atas E. Indirect hyperbilirubinemia because of eltrombopag in two cases after autologous hematopoietic stem cell transplantation. *J Cancer Res Ther.* 2023; 19: S970-S972.
92. Barisic S, Childs RW. Graft-versus-solid-tumor effect: From hematopoietic stem cell transplantation to adoptive cell therapies. *Stem Cells.* 2022; 40: 556-563.
93. Gandhi M, Sharma B, Nair S, Vaidya AD. Current insights into CAR T-cell-based therapies for myelodysplastic syndrome. *Pharm Res.* 2024; 41: 1757-1773.
94. Aboul-Soud MA, Alzahrani AJ, Mahmoud A. Induced pluripotent stem cells (iPSCs)-roles in regenerative therapies, disease modelling and drug screening. *Cells.* 2021; 10: 2319.
95. He R, Weng Z, Liu Y, Li B, Wang W, Meng W, et al. Application of induced pluripotent stem cells in malignant solid tumors. *Stem Cell Rev Rep.* 2023; 19: 2557-2575.
96. Zhang L, Tian L, Dai X, Yu H, Wang J, Lei A, et al. Pluripotent stem cell-derived CAR-macrophage cells with antigen-dependent anti-cancer cell functions. *J Hematol Oncol.* 2020; 13: 153.
97. Xue D, Lu S, Zhang H, Zhang L, Dai Z, Kaufman DS, et al. Induced pluripotent stem cell-derived engineered T cells, natural killer cells, macrophages, and dendritic cells in immunotherapy. *Trends Biotechnol.* 2023; 41: 907-922.
98. Kumar D, Gupta S, Gupta V, Tanwar R, Chandel A. Engineering the future of regenerative medicines in gut health with stem cell-derived intestinal organoids. *Stem Cell Rev Rep.* 2025; 21: 1449-1470.
99. Bayik D, Lathia JD. Cancer stem cell-immune cell crosstalk in tumour progression. *Nat Rev Cancer.* 2021; 21: 526-536.
100. Huang T, Song X, Xu D, Tiek D, Goenka A, Wu B, et al. Stem cell programs in cancer initiation, progression, and therapy resistance. *Theranostics.* 2020; 10: 8721-8743.
101. Wang C, Li Y, Jia L, Kim JK, Li J, Deng P, et al. CD276 expression enables squamous cell carcinoma stem cells to evade immune surveillance. *Cell Stem Cell.* 2021; 28: 1597-1613.e7.
102. Terzuoli E, Bellan C, Aversa S, Ciccone V, Morbidelli L, Giachetti A, et al. ALDH3A1 overexpression in melanoma and lung tumors drives cancer stem cell expansion, impairing immune surveillance through enhanced PD-L1 output. *Cancers.* 2019; 11: 1963.
103. Mortezaee K. Enriched cancer stem cells, dense stroma, and cold immunity: Interrelated events in pancreatic cancer. *J Biochem Mol Toxicol.* 2021; 35: e22708.
104. Rodríguez MM, Fiore E, Bayo J, Atorrasagasti C, García M, Onorato A, et al. 4Mu decreases CD47 expression on hepatic cancer stem cells and primes a potent antitumor T cell response induced by interleukin-12. *Mol Ther.* 2018; 26: 2738-2750.
105. Ho T, Msallam R. Tissues and tumor microenvironment (TME) in 3D: Models to shed light on immunosuppression in cancer. *Cells.* 2021; 10: 831.

106. Ali S, Borin TF, Piranlioglu R, Ara R, Lebedyeva I, Angara K, et al. Changes in the tumor microenvironment and outcome for TME-targeting therapy in glioblastoma: A pilot study. *PLoS One*. 2021; 16: e0246646.
107. Yang M, Li J, Gu P, Fan X. The application of nanoparticles in cancer immunotherapy: Targeting tumor microenvironment. *Bioact Mater*. 2021; 6: 1973-1987.
108. Hawkins ER, D'Souza RR, Klampatsa A. Armored CAR T-cells: The next chapter in T-cell cancer immunotherapy. *Biologics*. 2021; 15: 95-105.
109. Mashima H, Zhang R, Kobayashi T, Tsukamoto H, Liu T, Iwama T, et al. Improved safety of induced pluripotent stem cell-derived antigen-presenting cell-based cancer immunotherapy. *Mol Ther Methods Clin Dev*. 2021; 21: 171-179.
110. Wang LT, Liu KJ, Sytwu HK, Yen ML, Yen BL. Advances in mesenchymal stem cell therapy for immune and inflammatory diseases: Use of cell-free products and human pluripotent stem cell-derived mesenchymal stem cells. *Stem Cells Transl Med*. 2021; 10: 1288-1303.
111. Raturi M, Das K. Ethical issues facing hematopoietic stem cell transplantation in India. *Bull Cancer*. 2021; 108: 893-894.
112. Novoa JJ, Westra IM, Steeneveld E, Neves NF, Arendzen CH, Rajaei B, et al. Good manufacturing practice-compliant human induced pluripotent stem cells: From bench to putative clinical products. *Cytotherapy*. 2024; 26: 556-566.
113. Lappin T, Cheng T. An urgent need for standardization of stem cells and stem cell-derived products toward clinical applications. *Stem Cells Transl Med*. 2021; 10: S1-S3.
114. Chouaib B, Haack-Sørensen M, Chaubron F, Cuisinier F, Collart-Dutilleul PY. Towards the standardization of mesenchymal stem cell secretome-derived product manufacturing for tissue regeneration. *Int J Mol Sci*. 2023; 24: 12594.
115. Li YR, Dunn ZS, Yu Y, Li M, Wang P, Yang L. Advancing cell-based cancer immunotherapy through stem cell engineering. *Cell Stem Cell*. 2023; 30: 592-610.
116. Cheng C, Tang N, Li J, Cao S, Zhang T, Wei X, et al. Bacteria-free minicircle DNA system to generate integration-free CAR-T cells. *J Med Genet*. 2019; 56: 10-17.
117. Alqarni A, Jasim SA, Altalbawy FM, Kaur H, Kaur I, Rodriguez-Benites C, et al. Challenges and opportunities for cancer stem cell-targeted immunotherapies include immune checkpoint inhibitor, cancer stem cell-dendritic cell vaccine, chimeric antigen receptor immune cells, and modified exosomes. *J Biochem Mol Toxicol*. 2024; 38: e23719.
118. Zhen A, Carrillo MA, Mu W, Rezek V, Martin H, Hamid P, et al. Robust CAR-T memory formation and function via hematopoietic stem cell delivery. *PLoS Pathog*. 2021; 17: e1009404.
119. Kumar D, Tanwar R. Advancing NK cell therapies: Expansion and activation for next-generation immunotherapy. *Curr Cancer Ther Rev*. 2026; 22: 470-484.
1. Mim TJ, Ahmad I, Basher SR, Rahman MF, Ambati SR, Venkatagiri K, et al. Stem cells in organogenesis and regeneration. *Stem Cell Res Ther*. 2026; 17: 85.
2. Bryukhovetskiy I, Pak O, Khotimchenko Y, Bryukhovetskiy A, Sharma A, Sharma HS. Personalized therapy and stem cell transplantation for pro-inflammatory modulation of cancer stem cells microenvironment in glioblastoma. *Int Rev Neurobiol*. 2020; 151: 67-98.