

Review

**Diet and Prevention of Cardiovascular Disease**

Shanthi Mendis \*

Geneva Learning Foundation, Geneva, Switzerland; E-Mail: [prof.shanthi.mendis@gmail.com](mailto:prof.shanthi.mendis@gmail.com)\* **Correspondence:** Shanthi Mendis; E-Mail: [prof.shanthi.mendis@gmail.com](mailto:prof.shanthi.mendis@gmail.com)**Academic Editor:** Arrigo F. G. Cicero**Special Issue:** [Nutrition, Diet and Cardiovascular Disease](#)*Recent Progress in Nutrition*  
2024, volume 4, issue 3  
doi:10.21926/rpn.2403011**Received:** January 28, 2024  
**Accepted:** August 12, 2024  
**Published:** August 16, 2024**Abstract**

Promoting a healthy diet is an effective strategy for preventing cardiovascular disease (CVD). The characteristics of a healthy diet are known. However, an unhealthy diet has become a significant contributor to the global burden of diseases, mainly due to its effect on CVD, diabetes, and cancer. A healthy diet is not accessible and affordable to most people worldwide. Marketing influences food choices and promotes unhealthy diets that contribute to obesity, CVD, diabetes mellitus, and cancer. Governments are responsible for transforming food systems to deliver healthy, sustainable, and affordable diets using coherent policies, regulations, and legislation portfolios. This review highlights the role of a healthy diet in preventing CVD. It summarises the current scientific evidence underpinning dietary recommendations and the strategies for translating them into action.

**Keywords**

Cardiovascular disease; cardiovascular risk; coronary heart disease; stroke; healthy diet; dietary fats; dietary fibre; meat consumption; salt intake; sugar sweetened beverages; dietary recommendations; prevention; diabetes mellitus



© 2024 by the author. This is an open access article distributed under the conditions of the [Creative Commons by Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is correctly cited.

## **1. Introduction**

Cardiovascular Diseases (e.g., coronary heart disease (CHD), strokes, and heart failure) continue to account for the highest disease burden worldwide. In 2019, out of the 41 million deaths due to Non-Communicable Diseases (NCD), 17.9 million were due to Cardiovascular Diseases (CVD) [1]. Unhealthy diet, tobacco use, physical inactivity, and air pollution are the four major risk factors for NCD and CVD. They lead to cardiometabolic derangement, obesity, diabetes, hypertension, and hyperlipidemia), which drives up the CVD burden [2]. Reducing exposure to those described above behavioral, environmental, and metabolic risk factors is an effective strategy to prevent NCD and CVD [3] and significantly reduce high healthcare costs [2, 4]. Currently, many people are food insecure, malnourished, or overweight because they are unable to afford a healthy diet. In 2020, adult and childhood obesity increased alarmingly in all parts of the world, and an estimated 38.9 million children were overweight [5].

## **2. Role of Diet in Prevention of CVD**

Macro and micronutrients affect the development of CVD through multiple mechanisms, including their impact on blood lipids, blood pressure, blood clotting, blood glucose, body weight, and the gut microbiome [2, 6-13]. Guidelines recommend that a healthy diet include various foods while ensuring energy balance to avoid unhealthy weight gain [6-12]. Being overweight and obese increases the risk of NCD, including CVD, type 2 diabetes, and certain types of cancer [14]. Obesity is associated with higher all-cause mortality [15]. In addition to physical inactivity and genetic factors, macronutrient imbalance in the diet (e.g., the proportion of protein, fat, and carbohydrates) contributes to the development of overweight and obesity [14-17]. Another critical driver of obesity in all countries is the aggressive marketing of more processed, high-energy, and affordable food [17].

## **3. Healthy Diet**

A healthy diet should provide proteins, carbohydrates, fiber, fat, fluids, and micronutrients (minerals and vitamins) from a range of food groups [6-12] (Figure 1). Proteins from animal sources (milk, eggs, fish, and meat) fully complement the essential amino acids the body needs. Most plant-based foods do not contain all the essential amino acids. If the diet is entirely plant-based, it is necessary to consume a mixture of plant-based foods to obtain all the essential amino acids. Fruits and vegetables are rich sources of fiber and micronutrients. Most ultra-processed foods (e.g., ham, sausages, bacon, crisps, ice cream, biscuits) contain high saturated fat, salt, sugar, and food additives. Consumption of ultra-processed food has no health benefits and is associated with an increased risk of Non-Communicable Diseases and mental health disorders [13]. Herbs and spices (turmeric, saffron, black pepper, red pepper, ginger, garlic, onion, coriander, thyme, oregano, bay leaf, dill, sumac, cinnamon, cloves, cardamom, and many others) used in the preparation of food, are rich in phytonutrients and have been shown to have a range of beneficial health effects [18].


A Healthy Diet includes  
Fruits and vegetables, proteins  
carbohydrates, fat and fluids.

Fluids: 6-8 glasses /day  
(water, tea, coffee, avoid  
sweetened beverages)

On average, men need 2500 kcal  
and women need 2000 kcal per  
day. The exact amount of energy  
varies based on age, physical  
activity and health profile.

Healthy Eating Window  
Is less than 12hours/day  
(Consume the majority of  
caloric intake during lunch.  
Avoid late-night Snacking)

Portion sizes:  
<https://www.bda.uk.com/resource/food-facts-portion-sizes.html>



**Figure 1** Health promoting diet.

Based on scientific evidence, current dietary Guidelines recommend: i) restricting high-fat foods (e.g., butter and cream), replacing saturated fats (SF) with polyunsaturated (PUSF) and monounsaturated fats (MUSF), and avoiding trans fatty acids (TFA); ii) adequate intake of plant-based food from complex, high-fiber carbohydrates such as whole grains, legumes, fruits, vegetables, and nuts while restricting refined starches and free sugars; iii) reducing salt intake; iv) consumption of fish and lean meat in moderation but restricting red meat and processed meat; v) consumption of low-fat and/or fermented dairy products (e.g., yogurt); vi) reducing foods rich in cholesterol (e.g., eggs and shellfish) for those with diabetes or at higher risk of CVD and vii) restricting sugar-sweetened beverages.

Scientific evidence underpinning each of these recommendations is summarised below.

### **3.1 Restrict High Fat Foods, Substitute SF with PUSF and MUSF and Avoid TFA**

Dietary fat is the most energy-dense macronutrient. Foods and oils contain a mixture of fats and fatty acids: SF, PUSF, and MUSF. The primary sources of SF are fatty meat, dairy foods, lard, butter, ghee, palm oil, and coconut oil. There are two main families of PUSF: omega-3 and omega-6. PUSF and MUSF come mainly from plant sources: PUSF from nuts, soybean, safflower, sunflower, and sesame seeds, and fish; MUSF from avocados, olives, tree nuts (e.g., almonds, cashews, and pistachios) and seeds (flax and chia). Oils have a mixture of fats. Canola, corn, soybean, peanut, safflower, and sunflower oil are rich in PUSF. Olive oil is high in MUSF. Partial replacement of SF by PUSF and MUSF tends to lower total and low-density lipoprotein cholesterol (LDLC) levels [19].

Systematic reviews of randomized controlled trials (RCTs) find that reducing total fat intake reduces body weight [20, 21]. To reduce the risk of unhealthy weight gain, the WHO recommends that adults limit their total fat intake to 30% of their total energy intake or less. Reduced fat intake also results in a slight improvement in total cholesterol, LDL-C, and blood pressure. Meta-analysis

of four RCTs has shown that lowering SF and replacing it with PUSF lowers CHD by 29% [22-26]. This finding is supported by several other meta-analyses [27-31]. The reduction in CHD is similar to that achieved by statins [22]. Based on these and other findings [32-34], it is recommended that adults and children reduce SF intake to at least 10% of total energy intake, replacing SF in the diet with PUSF or MUSF from plant sources [8].

Strong evidence exists that lowering TFA intake reduces LDLC and is associated with a reduced risk of CVD [35]. Industrially produced TFA can be found in baked and fried foods (e.g., cookies, biscuits, pies) and pre-packaged snacks. Partially hydrogenated cooking oils and fats also contain TFA. WHO recommends consuming 1% or less of total energy intake as TFA [8]. Industrially produced TFA have no known health benefits and must be removed from the food supply through legislation or regulatory action.

Coconut oil, which contains 82% of SF, has been shown to significantly increase LDLC [36, 37]. The LDL cholesterol-raising property of cholesterol is confirmed by a recent systematic review of 7 controlled trials [38]. The review also found no difference in raising LDLC between coconut oil and other oils high in SF, such as butter, beef fat, or palm oil. Guidelines advise against using coconut oil because coconut oil increases LDL-C, which increases the risk of CVD [22].

### **3.2 Adequate Intake of Plant-Based Food: Wholegrains, Pulses, Fruits, Vegetables and Nuts**

Carbohydrates (e.g., starches, sugar, and fiber) are found primarily in plant-based foods such as grains, cereals, pulses, fruits, and vegetables. Carbohydrates that are slowly digested, such as whole grains, pulses, fruits, and vegetables, are considered high quality and beneficial to health [39]. Rapidly digested carbohydrates (e.g., sugars, white bread, white rice) are regarded as low quality, and their restriction is recommended [10]. Many studies have found that a higher intake of whole grains, vegetables, fruits, and pulses was associated with a significantly reduced risk of disease and mortality [40-48]. This included a nearly 20% decrease in all-cause mortality, a 10–20% decrease in the risk of CVD, a greater than 20% decrease in the risk of type 2 diabetes, and a 16% reduction in cancer risk. A systematic review analyzed data from 95 studies and found a nonlinear dose-response relationship between fruit and vegetable intake of up to 800 g per day and CHD, stroke, and CVD [42]. Fruits and vegetables are rich in nutrients- fiber, vitamins (e.g., vitamin C and folate), minerals (e.g., potassium, iron, and magnesium), and dietary bioactive compounds (antioxidants, polyphenols, and carotenoids). These nutrients act synergistically through various biological mechanisms to improve vascular and immune function and to lower blood cholesterol, blood pressure, inflammation, and platelet aggregation [49-53].

Intake of fiber, fruit, and vegetables also influences the composition and diversity of the microbial population in the gut [54]. Certain dietary constituents are known to be metabolized by intestinal bacteria into metabolites that affect CVD or diabetes [55]. Some of the health-promoting effects associated with consumption of dietary fiber are mediated by butyrate-producing microbes [56]. There is also evidence that polyphenols (e.g., phenolic acids, flavonoids, and lignans) from fruits, vegetables, and cereals could alter gut microorganisms by inhibiting potential pathogenic organisms (e.g, *Helicobacter pylori*, *Staphylococcus* sp.) while promoting the growth of beneficial bacteria, including *Lactobacillus* and *Bifidobacteria* [57, 58].

It is recommended that carbohydrate intake should come mainly from whole grains, pulses, fruits, and vegetables. It is recommended that adults consume at least 400 gm of fruits and vegetables (2

servings of fruits and 3 servings of vegetables) per day, excluding starchy vegetables such as potatoes and yams [6]. Intake of whole grains, pulses, fruits, and vegetables is highly variable across and within populations and is generally inadequate globally [59-65]. Less than 20–30% of individuals in many low and middle-income countries meet WHO recommendations for vegetable and fruit consumption [66, 67].

A high consumption of nuts is associated with a 15% lower risk of CVD incidence and a 23% lower risk of CVD mortality [68]. Consumption of nuts positively impacts blood lipids, blood pressure, inflammation, and markers of glycemic control [69]. In addition to unsaturated fats, nuts are rich in protein, phytosterols, fiber, minerals, vitamins, and phenolic compounds. Dietary Guidelines recommend 30 grams of nuts for adults most days of the week [70].

### **3.3 Reduce Salt (Sodium) Intake**

High intake of sodium, is a well-established cause of elevated blood pressure, increased risk of CVD, and chronic kidney disease [71, 72]. Results of RCTs report that salt reduction lowers blood pressure in both hypertensive and normotensive people [73]. A dose-response relation has been demonstrated between the magnitude of blood pressure lowering and sodium reduction in the diet [74]. A decrease of 4.4 g/day salt is associated with a mean reduction of 4.18 mmHg for systolic blood pressure and 2.06 mmHg for diastolic blood pressure [75]. Salt reduction is one of the recommendations for the nonpharmacological treatment of hypertension, in addition to weight reduction, dietary potassium supplementation, and physical activity [76].

Reducing dietary sodium intake can prevent a large number of fatal and non-fatal heart attacks at a meager cost. Hence, a 30% relative reduction in the mean population's intake of salt/sodium by 2030 is one of the nine global NCD targets [2]. This target is attainable if several cost-effective (Best Buy) policies are rapidly implemented to lower population sodium intake [2-4, 9, 77]. They include reducing the sodium content in food products, front-of-pack labeling to help consumers choose food products with lower sodium content, mass media campaigns to educate consumers, and implementing public food procurement policies to reduce sodium content in the food served or sold. Stepwise population salt reduction programmes are essential in low and middle-income countries where resource constraints and lack of health insurance limit access to antihypertensive medicines for many people with hypertension.

In high-income countries, most of the salt in the diet is from processed, restaurant, and fast foods [78], while in most low and middle-income countries, the primary salt source is added during cooking or in sauces [79]. WHO recommends a maximum intake of <2000 mg/day sodium (<5 g/day salt) in adults [9, 80].

WHO is monitoring the progress in implementing salt reduction policies in Member States [9]. As of October 2022, 5% of countries (n = 9) have implemented at least two mandatory sodium reduction policies and all sodium-related Best Buy policies for addressing NCD. 22% of countries (n = 43) have implemented at least one mandatory policy. One-third of the remaining Member States (n = 64) have implemented at least one voluntary policy to reduce sodium intake, while 29% (n = 56) have made a policy commitment towards sodium reduction.

Using low-sodium salt substitutes is increasingly considered a potential blood pressure-lowering strategy by national health authorities [81]. WHO is in the process of finalizing recommendations on the use of low-sodium salt substitutes.

### **3.4 Consumption of Fish and Lean Meat and Restriction of Red Meat and Processed Meat**

Fish such as mackerel, salmon, and sardine are good sources of omega-3 fat. Antioxidant and anti-inflammatory properties of omega-3 fats lower the risk of CVD by reducing blood clotting, protecting arteries from hardening, and reducing the level of triglycerides in the blood [82-84]. The results of a systematic review of 25 prospective cohort studies indicate that fish consumption is inversely associated with CVD mortality. The CVD mortality risk was decreased by 4% with an increase of 20 g of fish intake [84].

Meats are broadly categorized into red (beef, pork, lamb), white (chicken, turkey, and rabbit), and processed meat (sausages, bacon, and salami). Unprocessed and processed red meat consumption are both associated with a higher risk of CVD and diabetes [85, 86]. Currently, evidence does not suggest a beneficial or detrimental role of white meat consumption in the development of CVD [87].

### **3.5 Consumption of Low-Fat and/or Fermented Dairy Products**

Meta-analysis of cohort studies with CVD hard endpoints and RCTs investigating the effect on major cardiovascular risk factors provide evidence that moderate dairy consumption (up to 200 g/day) has no detrimental impact on cardiovascular health [88, 89]. Further studies are needed to determine whether fermented dairy products have cardiovascular benefits compared to non-fermented dairy products [90].

### **3.6 Reduce Foods Rich in Cholesterol**

Dietary cholesterol has been suggested to increase the risk of CVD. However, studies included in systematic reviews have lacked the methodologic rigor to enable firm conclusions regarding the effects of dietary cholesterol on CVD risk [91]. Meat, eggs, shellfish, and full-fat dairy products are rich in cholesterol. Meat and full-fat dairy products also have high levels of SF. Eggs are high in cholesterol but low in SF: a sizeable whole egg (50 g) contains 244 mg of cholesterol in the egg yolk but only 1.2 g of SF [92]. Evidence on the impact of consumption of eggs on CVD from observational studies is mixed [93]. Studies report no association [94, 95], increased risk of CVD [96], or decreased risk of stroke [97]. Using data from observational studies, it is challenging to determine the impact of any individual food independently of a dietary pattern. The increased risk of CVD with egg consumption reported in the above studies may be related to dietary patterns rather than eggs. There is evidence from some observational studies suggesting that higher egg consumption could be associated with a higher risk of CVD in people with diabetes [98-100]. However, results from RCTs suggest that including 6 to 12 eggs per week, in a balanced diet consistent with cardiovascular health, has no adverse effect on major CVD risk factors in individuals with type 2 diabetes [101]. These findings suggest that eggs could be consumed in low to moderate amounts ( $\leq 1$  egg/d) as part of a healthy diet. Eggs contain all essential amino acids and offer a complete source of protein, minerals, and vitamins.

### **3.7 Restrict Sugar Sweetened Beverages (SSB)**

Prospective cohort studies have investigated the impact of SSB intake on the risk of CVD. For example, a meta-analysis of 27 longitudinal studies of medium to high methodological quality

showed that SSB intake increased the risk of obesity (RR = 1.17; 95% C.I. 1.10-1.25), CHD (RR = 1.15; 9% C.I. 1.06-1.25), stroke (RR = 1.10; 9% C.I. 1.01-1.19) and type 2 diabetes (RR = 1.20; 95% C.I. 1.13-1.28) in adults [102]. These findings are consistent with the results of several other meta-analyses [103-106]. A meta-analysis of six cohort studies found an 8% higher risk of CVD mortality and CVD incidence per one serving per day increment in SSB [107].

Probable biological mechanisms of the association between SSB and CVD risk include a rapid rise in blood glucose and insulin after SSB consumption, a high glycemic load that leads to weight gain, inflammation, insulin resistance, and metabolic syndrome [108-110], increased blood pressure [111] and increased synthesis hepatic triglycerides, all of which increase the risk of CVD [112].

Artificially sweetened beverages (ASB), such as low, no-calorie sweetened beverages, are potential SSB replacements. However, a meta-analysis of prospective cohort studies has shown that compared with those in the lowest group, the relative risk comparing extreme groups of ASB consumption was 1.10 (0.98-1.23) for CHD, 1.19 (1.09-1.29) for stroke, and 1.32 (1.15-1.52) for CVD events [104]. Another meta-analysis reported a J-shaped association between ASB intakes and all-cause or CVD mortality. Daily consumption of 1.5, 2, and 2.5 servings of ASB was associated with 4%, 8%, and 13% higher risks of all-cause mortality. These findings suggest that ASB is not necessarily a healthier alternative to SSB [113]. Future studies should further investigate the association between ASB intakes and cause-specific mortality.

Evidence shows that frequent 100% fruit juice intake may promote weight gain [114]. Based on the results of prospective cohort studies, 1 serving per day of 100% fruit juice was associated with a higher increase in body mass index among children.

#### **4. Association between Healthy Dietary Patterns, Food Groups and CVD**

Prospective cohort studies report that adherence to specific quantifiable eating patterns is consistently associated with a lower risk of CVD [115-117]. They are the healthy Mediterranean-style eating, healthy US-Style eating, healthy vegetarian eating [118], the alternative healthy eating index [119], the alternate Mediterranean diet score [120], and the Dietary Approaches to Stop Hypertension (DASH) diet [121, 122]. The Mediterranean diet is high in vegetables, fruits, grains, legumes, nuts, and virgin olive oil. It includes moderate fish and wine and is low in red, processed meat, and added sugars. The vegetarian diet is based on various fruits, vegetables, legumes, and whole grains. It does not include meat, poultry, or fish. The DASH diet focuses on vegetables, fruits, whole grains, legumes, fat-free or low-fat dairy, and nuts and limits the intake of salt, cholesterol, total and SF, red and processed meats, sweets, and added sugars, including SSB. All these dietary patterns, including food groups associated with lower risk of CVD, have reduced all-cause deaths by 8-22% [120, 123], 19-28% CVD death, and 11-23% cancer death [124-126].

Results of a dose-response meta-analysis quantify the relation between the intake of 12 major food groups and the risk of CHD, stroke, and heart failure [127]. An inverse association was present for whole grains (RR<sub>CHD</sub>: 0.95 (95% CI: 0.92-0.98), vegetables and fruits (RR<sub>CHD</sub>: 0.97 (0.96-0.99), nuts (RR<sub>CHD</sub>: 0.67 (0.43-1.05), and fish consumption (RR<sub>CHD</sub>: 0.88 (0.79-0.99)). A positive association was present for red meat (RR<sub>CHD</sub>: 1.15 (1.08-1.23), processed meat (RR<sub>CHD</sub>: 1.27 (1.09-1.49), and SSB consumption (RR<sub>CHD</sub>: 1.17 (1.11-1.23)).

There is considerable evidence that specific dietary patterns can reduce cardiovascular events and mortality (Table 1). They include a Mediterranean diet pattern, plant-based diets, and low-fat diets.

**Table 1** Impact of diet on cardiovascular morbidity and mortality.

Type of diet	Impact	Evidence
Mediterranean diet	Effective primary and secondary prevention significantly reduces vascular events (myocardial infarction and stroke) and cardiovascular deaths.	[128] [129] [130]
Plant-based diet	Greater adherence to an overall plant-based dietary pattern is significantly associated with a lower risk of cardiovascular mortality and a lower risk of CVD incidence.	[131]
Low-fat diet	Reducing saturated fat intake for at least two years causes a potentially significant reduction in combined cardiovascular events. Greater reduction in saturated fat causes more substantial reductions in cardiovascular events.	[20]
	Low-fat diets reduce all-cause mortality and non-fatal myocardial infarction in patients with increased cardiovascular risk.	[132]

## 5. Nutrition of Infants and Children

Breastfeeding of infants and young children (aged 12–36 months) significantly impacts their survival, health, and development. The positive effects of breastfeeding are sustained over the life course [133]. The findings of a multi-country study demonstrate how inappropriate marketing undermines confidence in breastfeeding, posing a significant threat to the health of infants and children [134]. Across all countries studied, formula milk companies market their products using diverse tactics, including engaging women through online and offline channels, exploiting the anxieties and concerns of parents, distorting science to legitimize their products, and encouraging health professionals to promote formula milk products. Governments need to recognize this threat and strengthen national measures to prevent formula milk marketing, including through domestic legislation and accountability mechanisms.

Healthy eating behaviors are formed during childhood. Adopting healthy behaviors in childhood can be beneficial for preventing CVD in later life. Children are exposed to aggressive marketing of foods and non-alcoholic beverages that are high in SF, TFA, free sugars and/or salt (HFSS), which is harmful to health. WHO has released a new policy guideline to protect children from the detrimental impact of food marketing. The guideline recommends that countries implement mandatory policies to protect children from marketing HFSS foods and non-alcoholic beverages. The new recommendations are based on recent evidence on how exposure to food marketing affects children’s health, eating behaviors, and food choices [135]. The rise in the consumption of unhealthy food commodities in children is mainly driven by the aggressive marketing of transnational corporations [136, 137]. Regrettably, only a handful of governments have approved a policy framework to restrict all marketing to children up to 18 years of age of unhealthy foods and non-alcoholic beverages.



## **6. Implementation of Nutrition Action in Countries**

In 2017, the global burden of disease study attributed 11 million deaths and 255 million DALYs to dietary risk factors [138]. The study found that high intake of sodium, low intake of whole grains, and low intake of fruits were the leading dietary risk factors for deaths and DALYs globally. WHO's Member States have endorsed global targets for improving nutrition and have committed to monitoring progress. These targets are vital for identifying priority areas for action and catalyzing global and local change. There are targets for maternal, infant, and child nutrition and breastfeeding. In addition to the target on reducing salt, three global NCD targets to be attained by 2030 are partly related to diet: A 25% relative reduction in the prevalence of raised blood pressure or to contain the prevalence of raised blood pressure and halt the rise in diabetes and obesity [2, 4].

Based on the information in WHO's Global Database on the Implementation of Nutrition Action (GINA) [139], 189 countries have national policies that include goals to promote healthy diets. A total of 182 countries include in their policies goals aligned to the global nutrition targets for 2025: 129 exclusive breastfeeding and 141 children overweight. Further, 165 of the 189 countries have also integrated goals for diet-related NCD targets: 92 for sodium/salt intake, 82 for high blood pressure, 132 for diabetes, and 152 for overweight and obesity in adults and adolescents. There is moderate progress in policies to promote healthy diets. Nutrition labeling is implemented in 122 countries. Countries are also taking action in the following areas: food reformulation 60 (in 40 countries, the focus is on sodium/salt reduction); trans-fat bans 26; and fiscal policies to promote healthy diets 38 out of which 30 are on SSB.

## **7. Transformation of Food Systems to Deliver Environmentally Sustainable Health Promoting Diets**

In addition to a health focus, diets must consider factors that determine the environmental sustainability of food production. They include greenhouse gas emissions, freshwater usage, land use, non-renewable energy consumption, eutrophication, particulate matter, food loss, and waste. A Lancet Commission presented a healthy diet for both people and the planet [140]. It emphasizes a plant-based diet with a more significant proportion of whole grains, fruits, vegetables, nuts, and legumes and a smaller proportion of meat and dairy products. A few countries have already incorporated sustainability into their national dietary guidelines by including recommendations such as a predominantly plant-based diet, reducing food waste, and reducing consumption of red and processed meat [141-144].

Cost is a critical factor that determines the adoption of a healthy diet. There is concern that a healthy diet may not be affordable to low-income households in most settings due to the higher costs associated with healthy cuts of meat and the enormous amount of fruits, vegetables, legumes, and nuts in such a diet [145, 146]. About three billion people cannot afford a healthy diet and consume foods high in unhealthy fats, sugars and/or salt, which have become cheaper and more widely available in recent years.

WHO recommends implementing cost-effective and scalable food systems that focus on improving the nutritional quality of food and creating healthier food environments [147]. They include:

- Levying taxes on unhealthy food and subsidizing healthy options;
- regulating the marketing of food and non-alcoholic beverages to children;

- ensuring clear, informative, and accurate front-of-pack nutrition labels;
- reducing salt, sugars, and harmful fats in processed food;
- fortifying staple foods by adding extra vitamins and minerals;
- ensuring healthier diets are served or sold in public places;
- achieving coherence between trade and nutrition policies;
- and improving and supervising national food control systems.

Taxes need to be levied on unhealthy food options (such as foods high in fats, sugars and/or salt) to increase prices and discourage purchase and consumption. At the same time, subsidies can reduce the cost of healthy options (e.g., nuts, fruits, and vegetables) and encourage consumption.

Dietary variability- the availability of a wide variety of highly palatable energy-dense food- can compromise satiety learning, one of the critical cognitive determinants of food intake. Research shows that animals learn about their foods' energy content or satiating quality and adjust their intake to balance energy needs. Further research is needed to better define aspects in the modern food environment that undermine human satiety-learning, such as dietary variability [147].

In conclusion, compelling evidence defines a dietary pattern that is effective for preventing CVD and promoting health. People can benefit from it only if governments address physical, economic, and commercial determinants of the nexus between food and health. Further, government regulations must transform food systems to make them sustainable and allow the population access to affordable, healthy diets.

## **Abbreviations**

CHD	Coronary Heart Disease
CVD	Cardio Vascular Disease
DALYS	Disability Adjusted Life Years
DASH	Dietary Approaches to Stop Hypertension
LDLC	Low Density Lipoprotein Cholesterol
MUSF	Monounsaturated Fats
NCD	Non-Communicable Disease
PUSF	Polyunsaturated Fat
RCT	Randomized Controlled Trial
SF	Saturated Fat
SSB	Sugar Sweetened Beverages
TFA	Trans Fatty Acids
WHO	World Health Organization

## **Author Contributions**

SM was responsible for conceptualization, writing, review and editing of the paper.

## **Competing Interests**

Author was former Senior Adviser, Noncommunicable Diseases, World Health Organization, Geneva Switzerland.

## References

1. World Health Organization. World Health Statistics 2023: Monitoring Health for the SDGs, sustainable development goals. Geneva, Switzerland: World Health Organization; 2023.
2. World Health Organization. Global status report on noncommunicable diseases 2014. Geneva, Switzerland: World Health Organization; 2014.
3. Mendis S, Graham I, Narula J. Editorial: Reducing cardiovascular disease mortality and morbidity: Implementing cost-effective and sustainable preventive interventions. *Front Cardiovasc Med.* 2023; 10: 1236210.
4. World Health Organization. Implementation roadmap 2023-2030 for the Global Action Plan for the prevention and control of noncommunicable diseases 2013-2030. Geneva, Switzerland: World Health Organization; 2023.
5. FAO/IFAD/UNICEF/WFP/WHO. The state of food security and nutrition in the world 2021. Transforming food systems for food security, improved nutrition and affordable healthy diets for all. Rome, Italy: Food and Agriculture Organization of the United Nations; 2021.
6. World Health Organization. Carbohydrate intake for adults and children: WHO guideline. Geneva, Switzerland: World Health Organization; 2023.
7. World Health Organization. Total fat intake for the prevention of unhealthy weight gain in adults and children: WHO guideline. Geneva, Switzerland: World Health Organization; 2023.
8. World Health Organization. Saturated fatty acid and trans-fatty acid intake for adults and children: WHO guideline. Geneva, Switzerland: World Health Organization; 2023.
9. World Health Organization. WHO global report on sodium intake reduction. Geneva, Switzerland: World Health Organization; 2023.
10. World Health Organization. Guideline: Sugars intake for adults and children. Geneva, Switzerland: World Health Organization; 2015.
11. World Health Organization. Diet, nutrition and the prevention of chronic diseases. Report of a Joint WHO/FAO Expert Consultation. Geneva, Switzerland: World Health Organization; 2003; No. 916.
12. U.K. Government. The Eatwell Guide [Internet]. London, UK: U.K. Government; 2024. Available from: <https://www.gov.uk/government/publications/the-eatwell-guide>.
13. Dai S, Wellens J, Yang N, Li D, Wang J, Wang L, et al. Ultra-processed foods and human health: An umbrella review and updated meta-analyses of observational evidence. *Clin Nutr.* 2024; 43: 1386-1394.
14. World Health Organization. Obesity and overweight—Key facts [Internet]. Geneva, Switzerland: World Health Organization; 2021. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.
15. Di Angelantonio E, Bhupathiraju SN, Wormser D, Gao P, Kaptoge S, De Gonzalez AB, et al. Body-mass index and all-cause mortality: Individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet.* 2016; 388: 776-786.
16. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, et al. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med.* 2017; 377: 13-27.
17. Chavez-Ugalde Y, Jago R, Toumpakari Z, Egan M, Cummins S, White M, et al. Conceptualizing the commercial determinants of dietary behaviors associated with obesity: A systematic review using principles from critical interpretative synthesis. *Obes Sci Pract.* 2021; 7: 473-486.

18. Drewnowski A. A novel Nutrient Rich Food (NRFa11.3) score uses flavonoids and carotenoids to identify antioxidant-rich spices, herbs, vegetables, and fruit. *Front Nutr.* 2024; 11: 1386328.
19. Froyen E, Burns-Whitmore B. The effects of linoleic acid consumption on lipid risk markers for cardiovascular disease in healthy individuals: A review of human intervention trials. *Nutrients.* 2020; 12: 2329.
20. Hooper L, Abdelhamid AS, Jimoh OF, Bunn D, Skeaff C. Effects of total fat intake on body fatness in adults. *Cochrane Database Syst Rev.* 2020. doi: 10.1002/14651858.CD013636.
21. Naude CE, Visser ME, Nguyen KA, Durao S, Schoonees A. Effects of total fat intake on bodyweight in children. *Cochrane Database Syst Rev.* 2018. doi: 10.1002/14651858.CD012960.pub2.
22. Sacks FM, Lichtenstein AH, Wu JH, Appel LJ, Creager MA, Kris-Etherton PM, et al. Dietary fats and cardiovascular disease: A presidential advisory from the American Heart Association. *Circulation.* 2017; 136: e1-e23. Erratum in: *Circulation.* 2017; 136: e195.
23. Dayton S, Pearce ML, Hashimoto S, Dixon WJ, Tomiyasu U. A controlled clinical trial of a diet high in unsaturated fat in preventing complications of atherosclerosis. *Circulation.* 1969; 40: II-1-II-63.
24. Leren P. The Oslo diet-heart study: Eleven-year report. *Circulation.* 1970; 42: 935-942.
25. Controlled trial of soya-bean oil in myocardial infarction. *Lancet.* 1968; 2: 693-699.
26. Turpeinen O, Karvonen MJ, Pekkarinen M, Miettinen M, Elosuo R, Paavilainen E. Dietary prevention of coronary heart disease: The Finnish mental hospital study. *Int J Epidemiol.* 1979; 8: 99-118.
27. Miettinen M, Turpeinen O, Karvonen MJ, Pekkarinen M, Paavilainen E, Elosuo R. Dietary prevention of coronary heart disease in women: The Finnish Mental Hospital Study. *Int J Epidemiol.* 1983; 12: 17-25.
28. Luo S, Hou H, Wang Y, Li Y, Zhang L, Zhang H, et al. Effects of omega-3, omega-6, and total dietary polyunsaturated fatty acid supplementation in patients with atherosclerotic cardiovascular disease: A systematic review and meta-analysis. *Food Funct.* 2024; 15: 1208-1222.
29. Chowdhury R, Warnakula S, Kunutsor S, Crowe F, Ward HA, Johnson L, et al. Association of dietary, circulating, and supplement fatty acids with coronary risk: A systematic review and meta-analysis. *Ann Intern Med.* 2014; 160: 398-406.
30. Mozaffarian D, Micha R, Wallace S. Effects on coronary heart disease of increasing polyunsaturated fat in place of saturated fat: A systematic review and meta-analysis of randomized controlled trials. *PLoS Med.* 2010; 7: e1000252.
31. Hooper L, Martin N, Abdelhamid A, Davey Smith G. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database Syst Rev.* 2015; CD011737. doi: 10.1002/14651858.CD011737.
32. Farvid MS, Ding M, Pan A, Sun Q, Chiuve SE, Steffen LM, et al. Dietary linoleic acid and risk of coronary heart disease: A systematic review and meta-analysis of prospective cohort studies. *Circulation.* 2014; 130: 1568-1578.
33. Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease. *Am J Clin Nutr.* 2010; 91: 535-546.

34. Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: A meta-analysis of 60 controlled trials. *Am J Clin Nutr.* 2003; 77: 1146-1155.
35. Bendsen NT, Christensen R, Bartels EM, Astrup A. Consumption of industrial and ruminant trans fatty acids and risk of coronary heart disease: A systematic review and meta-analysis of cohort studies. *Eur J Clin Nutr.* 2011; 65: 773-783.
36. Jeyakumar SM, Damayanti K, Rajkumar Ponday L, Acharya V, Koppala SR, Putcha UK, et al. Assessment of virgin coconut oil in a balanced diet on indicators of cardiovascular health in non-obese volunteers: A human metabolic study. *Diabetes Metab Syndr.* 2023; 17: 102844.
37. Voon PT, Ng TK, Lee VK, Nesaretnam K. Diets high in palmitic acid (16:0), lauric and myristic acids (12:0 + 14:0), or oleic acid (18:1) do not alter postprandial or fasting plasma homocysteine and inflammatory markers in healthy Malaysian adults. *Am J Clin Nutr.* 2011; 94: 1451-1457.
38. Eyres L, Eyres MF, Chisholm A, Brown RC. Coconut oil consumption and cardiovascular risk factors in humans. *Nutr Rev.* 2016; 74: 267-280.
39. Reynolds A, Mann J, Cummings J, Winter N, Mete E, Te Morenga L. Carbohydrate quality and human health: A series of systematic reviews and meta-analyses. *Lancet.* 2019; 393: 434-445.
40. The InterAct Consortium. Dietary fibre and incidence of type 2 diabetes in eight European countries: The EPIC-InterAct Study and a meta-analysis of prospective studies. *Diabetologia.* 2015; 58: 1394-1408.
41. Aune D, Keum N, Giovannucci E, Fadnes LT, Boffetta P, Greenwood DC, et al. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: Systematic review and dose–response meta-analysis of prospective studies. *BMJ.* 2016; 353: i2716.
42. Aune D, Giovannucci E, Boffetta P, Fadnes LT, Keum N, Norat T, et al. Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality: A systematic review and dose–response meta-analysis of prospective studies. *Int J Epidemiol.* 2017; 46: 1029-1056.
43. Viguiiyouk E, Glenn AJ, Nishi SK, Chiavaroli L, Seider M, Khan T, et al. Associations between dietary pulses alone or with other legumes and cardiometabolic disease outcomes: An umbrella review and updated systematic review and meta-analysis of prospective cohort studies. *Adv Nutr.* 2019; 10: S308-S319.
44. Reynolds AN, Diep Pham HT, Montez J, Mann J. Dietary fibre intake in childhood or adolescence and subsequent health outcomes: A systematic review of prospective observational studies. *Diabetes Obes Metab.* 2020; 22: 2460-2467.
45. Mytton OT, Nnoaham K, Eyles H, Scarborough P, Ni Mhurchu C. Systematic review and meta-analysis of the effect of increased vegetable and fruit consumption on body weight and energy intake. *BMC Public Health.* 2014; 14: 886.
46. Mytton OT, Nnoaham K, Eyles H, Scarborough P, Ni Mhurchu C. Erratum to: Systematic review and meta-analysis of the effect of increased vegetable and fruit consumption on body weight and energy intake. *BMC Public Health.* 2017; 17: 662.
47. Afshin A, Micha R, Khatibzadeh S, Mozaffarian D. Consumption of nuts and legumes and risk of incident ischemic heart disease, stroke, and diabetes: A systematic review and meta-analysis. *Am J Clin Nutr.* 2014; 100: 278-288.

48. Marventano S, Izquierdo Pulido M, Sánchez-González C, Godos J, Speciani A, Galvano F, et al. Legume consumption and CVD risk: A systematic review and meta-analysis. *Public Health Nutr.* 2017; 20: 245-254.
49. Anderson JW, Baird P, Davis Jr RH, Ferreri S, Knudtson M, Koraym A, et al. Health benefits of dietary fiber. *Nutr Rev.* 2009; 67: 188-205.
50. Boeing H, Bechthold A, Bub A, Ellinger S, Haller D, Kroke A, et al. Critical review: Vegetables and fruit in the prevention of chronic diseases. *Eur J Nutr.* 2012; 51: 637-663.
51. Broekmans WM, Klöpping-Ketelaars IA, Schuurman CR, Verhagen H, van den Berg H, Kok FJ, et al. Fruits and vegetables increase plasma carotenoids and vitamins and decrease homocysteine in humans. *J Nutr.* 2000; 130: 1578-1583.
52. Macready AL, George TW, Chong MF, Alimbetov DS, Jin Y, Vidal A, et al. Flavonoid-rich fruit and vegetables improve microvascular reactivity and inflammatory status in men at risk of cardiovascular disease—FLAVURS: A randomized controlled trial. *Am J Clin Nutr.* 2014; 99: 479-489.
53. Threapleton DE, Greenwood DC, Evans CE, Cleghorn CL, Nykjaer C, Woodhead C, et al. Dietary fiber intake and risk of first stroke: A systematic review and meta-analysis. *Stroke.* 2013; 44: 1360-1368.
54. Yang Q, Liang Q, Balakrishnan B, Belobrajdic DP, Feng QJ, Zhang W. Role of dietary nutrients in the modulation of gut microbiota: A narrative review. *Nutrients.* 2020; 12: 381.
55. Yang Y, Karampoor S, Mirzaei R, Borozdkin L, Zhu P. The interplay between microbial metabolites and macrophages in cardiovascular diseases: A comprehensive review. *Int Immunopharmacol.* 2023; 121: 110546.
56. Björkegren JL, Lusis AJ. Atherosclerosis: Recent developments. *Cell.* 2022; 185: 1630-1645.
57. Selma MV, Espin JC, Tomas-Barberan FA. Interaction between phenolics and gut microbiota: Role in human health. *J Agric Food Chem.* 2009; 57: 6485-6501.
58. Parkar SG, Stevenson DE, Skinner MA. The potential influence of fruit polyphenols on colonic microflora and human gut health. *Int J Food Microbiol.* 2008; 124: 295-298.
59. Micha R, Khatibzadeh S, Shi P, Andrews KG, Engell RE, Mozaffarian D. Global, regional and national consumption of major food groups in 1990 and 2010: A systematic analysis including 266 country specific nutrition surveys worldwide. *BMJ Open.* 2015; 5: e008705.
60. Meynier A, Chanson-Rollé A, Riou E. Main factors influencing whole grain consumption in children and adults: A narrative review. *Nutrients.* 2020; 12: 2217.
61. Miller KB. Review of whole grain and dietary fiber recommendations and intake levels in different countries. *Nutr Rev.* 2020; 78: 29-36.
62. Stephen AM, Champ MM, Cloran SJ, Fleith M, van Lieshout L, Mejbourn H, et al. Dietary fibre in Europe: Current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev.* 2017; 30: 149-190.
63. Food and Agriculture Organization of the United Nations. The global economy of pulses [Internet]. Rome, Italy: Food and Agriculture Organization of the United Nations; 2019. Available from: <https://www.fao.org/documents/card/en/c/i7108en>.
64. McGill CR, Fulgoni III VL, Devareddy L. Ten-year trends in fiber and whole grain intakes and food sources for the United States population: National Health and Nutrition Examination Survey 2001–2010. *Nutrients.* 2015; 7: 1119-1130.

65. Miller V, Yusuf S, Chow CK, Dehghan M, Corsi DJ, Lock K, et al. Availability, affordability, and consumption of fruits and vegetables in 18 countries across income levels: Findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet Glob Health*. 2016; 4: e695-e703.
66. Darfour-Oduro SA, Buchner DM, Andrade JE, Grigsby-Toussaint DS. A comparative study of fruit and vegetable consumption and physical activity among adolescents in 49 low-and-middle-income countries. *Sci Rep*. 2018; 8: 1623.
67. Frank SM, Webster J, McKenzie B, Geldsetzer P, Manne-Goehler J, Andall-Brereton G, et al. Consumption of fruits and vegetables among individuals 15 years and older in 28 low- and middleincome countries. *J Nutr*. 2019; 149: 1252-1259.
68. Becerra-Tomás N, Paz-Graniel I, WC Kendall C, Kahleova H, Rahelić D, Sievenpiper JL, et al. Nut consumption and incidence of cardiovascular diseases and cardiovascular disease mortality: A meta-analysis of prospective cohort studies. *Nutr Rev*. 2019; 77: 691-709.
69. Glenn AJ, Aune D, Freisling H, Mohammadifard N, Kendall CW, Salas-Salvadó J, et al. Nuts and cardiovascular disease outcomes: A review of the evidence and future directions. *Nutrients*. 2023; 15: 911.
70. National Health and Medical Research Council. Eat for health Australian dietary guidelines [Internet]. Canberra, Australia: NHMRC Publications; 2013. Available from: [https://www.eatforhealth.gov.au/sites/default/files/files/the\\_guidelines/n55\\_australian\\_dietary\\_guidelines.pdf](https://www.eatforhealth.gov.au/sites/default/files/files/the_guidelines/n55_australian_dietary_guidelines.pdf).
71. Mozaffarian D, Fahimi S, Singh GM, Micha R, Khatibzadeh S, Engell RE, et al. Global sodium consumption and death from cardiovascular causes. *N Engl J Med*. 2014; 371: 624-634.
72. Filippini T, Malavolti M, Whelton PK, Naska A, Orsini N, Vinceti M. Blood pressure effects of sodium reduction: Dose-response meta-analysis of experimental studies. *Circulation*. 2021; 143: 1542-1567.
73. He FJ, Tan M, Ma Y, MacGregor GA. Salt reduction to prevent hypertension and cardiovascular disease: JACC state-of-the-art review. *J Am Coll Cardiol*. 2020; 75: 632-647.
74. Huang L, Trieu K, Yoshimura S, Neal B, Woodward M, Campbell NR, et al. Effect of dose and duration of reduction in dietary sodium on blood pressure levels: Systematic review and meta-analysis of randomised trials. *BMJ*. 2020; 368: m315.
75. He FJ, Li J, Macgregor GA. Effect of longer-term modest salt reduction on blood pressure. *Cochrane Database Syst Rev*. 2013. doi: 10.1002/14651858.CD004937.pub2.
76. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: A report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2018; 71: e127-e248.
77. World Health Organization. Advancing the global agenda on prevention and control of noncommunicable diseases 2000 to 2020: Looking forwards to 2030. Geneva, Switzerland: World Health Organization; 2023.
78. James WP, Ralph A, Sanchez-Castillo C. The dominance of salt in manufactured food in the sodium intake of affluent societies. *Lancet*. 1987; 329: 426-429.
79. Anderson CA, Appel LJ, Okuda N, Brown IJ, Chan Q, Zhao L, et al. Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: The INTERMAP study. *J Am Diet Assoc*. 2010; 110: 736-745.

80. World Health Organization. Guideline: Sodium intake for adults and children. Geneva, Switzerland: World Health Organization; 2012.
81. Brand A, Visser ME, Schoonees A, Naude CE. Replacing salt with low-sodium salt substitutes (LSS) for cardiovascular health in adults, children and pregnant women. *Cochrane Database Syst Rev*. 2022. doi: 10.1002/14651858.CD015207.
82. Elagizi A, Lavie CJ, O'Keefe E, Marshall K, O'Keefe JH, Milani RV. An update on omega-3 polyunsaturated fatty acids and cardiovascular health. *Nutrients*. 2021; 13: 204.
83. Djuricic I, Calder PC. Beneficial outcomes of omega-6 and omega-3 polyunsaturated fatty acids on human health: An update for 2021. *Nutrients*. 2021; 13: 2421.
84. Jiang L, Wang J, Xiong K, Xu L, Zhang B, Ma A. Intake of fish and marine n-3 polyunsaturated fatty acids and risk of cardiovascular disease mortality: A meta-analysis of prospective cohort studies. *Nutrients*. 2021; 13: 2342.
85. Shi W, Huang X, Schooling CM, Zhao JV. Red meat consumption, cardiovascular diseases, and diabetes: A systematic review and meta-analysis. *Eur Heart J*. 2023; 44: 2626-2635.
86. de Medeiros GC, Mesquita GX, Lima SC, Silva DF, de Azevedo KP, Pimenta ID, et al. Associations of the consumption of unprocessed red meat and processed meat with the incidence of cardiovascular disease and mortality, and the dose-response relationship: A systematic review and meta-analysis of cohort studies. *Crit Rev Food Sci Nutr*. 2023; 63: 8443-8456.
87. Ramel A, Nwaru BI, Lamberg-Allardt C, Thorisdottir B, Bärebring L, Söderlund F, et al. White meat consumption and risk of cardiovascular disease and type 2 diabetes: A systematic review and meta-analysis. *Food Nutr Res*. 2023. doi: 10.29219/fnr.v67.9543.
88. Giosuè A, Calabrese I, Vitale M, Riccardi G, Vaccaro O. Consumption of dairy foods and cardiovascular disease: A systematic review. *Nutrients*. 2022; 14: 831.
89. Guo J, Astrup A, Lovegrove JA, Gijsbers L, Givens DJ, Soedamah-Muthu SS. Milk and dairy consumption and risk of cardiovascular diseases and all-cause mortality: Dose-response meta-analysis of prospective cohort studies. *Eur J Epidemiol*. 2017; 32: 269-287.
90. Koskinen TT, Virtanen HEK, Voutilainen S, Tuomainen TP, Mursu J, Virtanen JK. Intake of fermented and non-fermented dairy products and risk of incident CHD: The Kuopio Ischaemic Heart Disease Risk Factor Study. *Br J Nutr*. 2018; 120: 1288-1297.
91. Berger S, Raman G, Vishwanathan R, Jacques PF, Johnson EJ. Dietary cholesterol and cardiovascular disease: A systematic review and meta-analysis. *Am J Clin Nutr*. 2015; 102: 276-294.
92. Food Standards Australia New Zealand. AUSNUT 2011–13 – Australian food composition database [Internet]. Canberra, Australia: Food Standards Australia New Zealand; 2014. Available from: <https://www.foodstandards.gov.au/science-data/food-composition-databases/ausnut-2011-13>.
93. Carter S, Connole ES, Hill AM, Buckley JD, Coates AM. Eggs and cardiovascular disease risk: An update of recent evidence. *Curr Atheroscler Rep*. 2023; 25: 373-380.
94. Godos J, Micek A, Brzostek T, Toledo E, Iacoviello L, Astrup A, et al. Egg consumption and cardiovascular risk: A dose-response meta-analysis of prospective cohort studies. *Eur J Nutr*. 2021; 60: 1833-1862.
95. Ma W, Zhang Y, Pan L, Wang S, Xie K, Deng S, et al. Association of egg consumption with risk of all-cause and cardiovascular disease mortality: A systematic review and dose-response meta-analysis of observational studies. *J Nutr*. 2022; 152: 2227-2237.



96. Yang PF, Wang CR, Hao FB, Peng Y, Wu JJ, Sun WP, et al. Egg consumption and risks of all-cause and cause-specific mortality: A dose–response meta-analysis of prospective cohort studies. *Nutr Rev.* 2022; 80: 1739-1754.
97. Marventano S, Godos J, Tieri M, Ghelfi F, Titta L, Lafranconi A, et al. Egg consumption and human health: An umbrella review of observational studies. *Int J Food Sci Nutr.* 2020; 71: 325-331.
98. Shin JY, Xun P, Nakamura Y, He K. Egg consumption in relation to risk of cardiovascular disease and diabetes: A systematic review and meta-analysis. *Am J Clin Nutr.* 2013; 98: 146-159.
99. Rong Y, Chen L, Zhu T, Song Y, Yu M, Shan Z, et al. Egg consumption and risk of coronary heart disease and stroke: Dose-response meta-analysis of prospective cohort studies. *BMJ.* 2013; 346: e8539.
100. Li MY, Chen JH, Chen C, Kang YN. Association between egg consumption and cholesterol concentration: A systematic review and meta-analysis of randomized controlled trials. *Nutrients.* 2020; 12: 1995.
101. Richard C, Cristall L, Fleming E, Lewis ED, Ricupero M, Jacobs RL, et al. Impact of egg consumption on cardiovascular risk factors in individuals with Type 2 diabetes and at risk for developing diabetes: A systematic review of randomized nutritional intervention studies. *Can J Diabetes.* 2017; 41: 453-463.
102. Santos LP, Gigante DP, Delpino FM, Maciel AP, Bielemann RM. Sugar sweetened beverages intake and risk of obesity and cardiometabolic diseases in longitudinal studies: A systematic review and meta-analysis with 1.5 million individuals. *Clin Nutr ESPEN.* 2022; 51: 128-142.
103. Li B, Yan N, Jiang H, Cui M, Wu M, Wang L, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages and fruit juices and risk of type 2 diabetes, hypertension, cardiovascular disease, and mortality: A meta-analysis. *Front Nutr.* 2023; 10: 1019534.
104. Jiang YW, Zhang YB, Pan A. Consumption of sugar-sweetened beverages and artificially sweetened beverages and risk of cardiovascular disease: A meta-analysis. *Chin J Prev Med.* 2021; 55: 1159-1167.
105. Yang B, Glenn AJ, Liu Q, Madsen T, Allison MA, Shikany JM, et al. Added sugar, sugar-sweetened beverages, and artificially sweetened beverages and risk of cardiovascular disease: Findings from the women's health initiative and a network meta-analysis of prospective studies. *Nutrients.* 2022; 14: 4226.
106. Meng Y, Li S, Khan J, Dai Z, Li C, Hu X, et al. Sugar- and artificially sweetened beverages consumption linked to type 2 diabetes, cardiovascular diseases, and all-cause mortality: A systematic review and dose-response meta-analysis of prospective cohort studies. *Nutrients.* 2021; 13: 2636.
107. Yin J, Zhu Y, Malik V, Li X, Peng X, Zhang FF, et al. Intake of sugar-sweetened and low-calorie sweetened beverages and risk of cardiovascular disease: A meta-analysis and systematic review. *Adv Nutr.* 2021; 12: 89-101.
108. Pacheco LS, Tobias DK, Li Y, Bhupathiraju SN, Willett WC, Ludwig DS, et al. Sugar-sweetened or artificially-sweetened beverage consumption, physical activity, and risk of cardiovascular disease in adults: A prospective cohort study. *Am J Clin Nutr.* 2024; 119: 669-681.
109. Ludwig DS. The glycemic index: Physiological mechanisms relating to obesity, diabetes, and cardiovascular disease. *JAMA.* 2002; 287: 2414-2423.

110. Malik VS, Hu FB. Sugar-sweetened beverages and cardiometabolic health: An update of the evidence. *Nutrients*. 2019; 11: 1840.
111. Brown IJ, Stamler J, Van Horn L, Robertson CE, Chan Q, Dyer AR, et al. Sugar-sweetened beverage, sugar intake of individuals, and their blood pressure: International study of macro/micronutrients and blood pressure. *Hypertension*. 2011; 57: 695-701.
112. Fried SK, Rao SP. Sugars, hypertriglyceridemia, and cardiovascular disease. *Am J Clin Nutr*. 2003; 78: 873S-880S.
113. Zhang YB, Jiang YW, Chen JX, Xia PF, Pan A. Association of consumption of sugar-sweetened beverages or artificially sweetened beverages with mortality: A systematic review and dose-response meta-analysis of prospective cohort studies. *Adv Nutr*. 2021; 12: 374-383.
114. Nguyen M, Jarvis SE, Chiavaroli L, Mejia SB, Zurbau A, Khan TA, et al. Consumption of 100% fruit juice and body weight in children and adults: A systematic review and meta-analysis. *JAMA Pediatr*. 2024; 178: 237-246.
115. Shan Z, Li Y, Baden MY, Bhupathiraju SN, Wang DD, Sun Q, Rexrode KM, et al. Association between healthy eating patterns and risk of cardiovascular disease. *JAMA Intern Med*. 2020; 180: 1090-1100.
116. Nestel PJ, Mori TA. Dietary patterns, dietary nutrients and cardiovascular disease. *Rev Cardiovasc Med*. 2022; 23: 17.
117. Chen W, Zhang S, Hu X, Chen F, Li D. A review of healthy dietary choices for cardiovascular disease: From individual nutrients and foods to dietary patterns. *Nutrients*. 2023; 15: 4898.
118. U.S. Department of Health and Human Services and U.S. Department of Agriculture. 2015 – 2020 Dietary Guidelines for Americans [Internet]. Rockville, MD: U.S. Department of Health and Human Services and U.S. Department of Agriculture; 2015. Available from: <http://health.gov/dietaryguidelines/2015/guidelines/>.
119. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, et al. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr*. 2012; 142: 1009-1018.
120. Fung TT, Rexrode KM, Mantzoros CS, Manson JE, Willett WC, Hu FB. Mediterranean diet and incidence of and mortality from coronary heart disease and stroke in women. *Circulation*. 2009; 119: 1093-1100. Erratum in: *Circulation*. 2009; 119: e379.
121. Filippou CD, Tsioufis CP, Thomopoulos CG, Mihas CC, Dimitriadis KS, Sotiropoulou LI, et al. Dietary approaches to stop hypertension (DASH) diet and blood pressure reduction in adults with and without hypertension: A systematic review and meta-analysis of randomized controlled trials. *Adv Nutr*. 2020; 11: 1150-1160.
122. Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med*. 2008; 168: 713-720.
123. Schwingshackl L, Hoffmann G. Diet quality as assessed by the healthy eating index, the alternate healthy eating index, the dietary approaches to stop hypertension score, and health outcomes: A systematic review and meta-analysis of cohort studies. *J Acad Nutr Diet*. 2015; 115: 780-800.e5.
124. George SM, Ballard-Barbash R, Manson JE, Reedy J, Shikany JM, Subar AF, et al. Comparing Indices of diet quality with chronic disease mortality risk in postmenopausal women in the women's health initiative observational study: Evidence to inform national dietary guidance. *Am J Epidemiol*. 2014; 180: 616-625.

125. Liese AD, Krebs-Smith SM, Subar AF, George SM, Harmon BE, Neuhauser ML, et al. The dietary patterns methods project: Synthesis of findings across cohorts and relevance to dietary guidance. *J Nutr.* 2015; 145: 393-402.
126. Reedy J, Krebs-Smith SM, Miller PE, Liese AD, Kahle LL, Park Y, et al. Higher diet quality is associated with decreased risk of all-cause, cardiovascular disease, and cancer mortality among older adults. *J Nutr.* 2014; 144: 881-889.
127. Bechthold A, Boeing H, Schwedhelm C, Hoffmann G, Knüppel S, Iqbal K, et al. Food groups and risk of coronary heart disease, stroke and heart failure: A systematic review and dose-response meta-analysis of prospective studies. *Crit Rev Food Sci Nutr.* 2019; 59: 1071-1090.
128. Sebastian SA, Padda I, Johal G. Long-term impact of mediterranean diet on cardiovascular disease prevention: A systematic review and meta-analysis of randomized controlled trials. *Curr Probl Cardiol.* 2024; 49: 102509.
129. Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, et al. Primary prevention of cardiovascular disease with a Mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med.* 2018; 378: e34.
130. Doundoulakis I, Farmakis IT, Theodoridis X, Konstantelos A, Christoglou M, Kotzakioulafi E, et al. Effects of dietary interventions on cardiovascular outcomes: a network meta-analysis. *Nutr Rev.* 2024; 82: 715-725.
131. Quek J, Lim G, Lim WH, Ng CH, So WZ, Toh J, et al. The Association of Plant-Based Diet With Cardiovascular Disease and Mortality: A Meta-Analysis and Systematic Review of Prospect Cohort Studies. *Front Cardiovasc Med.* 2021; 8: 756810.
132. Karam G, Agarwal A, Sadeghirad B, Jalink M, Hitchcock CL, Ge L, et al. Comparison of seven popular structured dietary programmes and risk of mortality and major cardiovascular events in patients at increased cardiovascular risk: Systematic review and network meta-analysis. *BMJ* 2023; 380: e072003.
133. Horta BL, Victora CG. Long-term effects of breastfeeding: A systematic review. Geneva, Switzerland: World Health Organization; 2013.
134. World Health Organization and the United Nations Children's Fund. How the marketing of formula milk influences our decisions on infant feeding. Geneva, Switzerland: World Health Organization and the United Nations Children's Fund; 2022.
135. World Health Organization. Policies to protect children from the harmful impact of food marketing: WHO guideline. Geneva, Switzerland: World Health Organization; 2023.
136. Moodie R, Stuckler D, Monteiro C, Sheron N, Neal B, Thamarangsi T, et al. Profits and pandemics: Prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *Lancet.* 2013; 381: 670-679.
137. Wiist WH. The corporate play book, health, and democracy: The snack food and beverage industry's tactics in context. In: *Sick societies: Responding to the global challenge of chronic disease.* Oxford: Oxford University Press; 2012. pp. 204-216.
138. Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, et al. Health effects of dietary risks in 195 countries, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2019; 393: 1958-1972.
139. World Health Organization. Global database on the Implementation of Nutrition Action (GINA) [Internet]. Geneva, Switzerland: World Health Organization; 2024. Available from: <https://extranet.who.int/nutrition/gina/en/home>.

140. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the anthropocene: The EAT–Lancet commission on healthy diets from sustainable food systems. *Lancet*. 2019; 393: 447-492.
141. Batis C, Marrón-Ponce JA, Stern D, Vandevijvere S, Barquera S, Rivera JA. Adoption of healthy and sustainable diets in Mexico does not imply higher expenditure on food. *Nat Food*. 2021; 2: 792-801.
142. Goulding T, Lindberg R, Russell CG. The affordability of a healthy and sustainable diet: An Australian case study. *Nutr J*. 2020; 19: 109.
143. Swedish National Food Agency. Find your way to eat greener, not too much and be active. Uppsala, Sweden: National Food Agency; 2015.
144. Serra-Majem L, Tomaino L, Dernini S, Berry EM, Lairon D, Ngo de la Cruz J, et al. Updating the mediterranean diet pyramid towards sustainability: Focus on environmental concerns. *Int J Environ Res Public Health*. 2020; 17: 8758.
145. Barosh L, Friel S, Engelhardt K, Chan L. The cost of a healthy and sustainable diet--who can afford it? *Aust N Z J Public Health*. 2014; 38: 7-12.
146. WHO. Food Systems Delivering Better Health. A new narrative to guide policy and practice for better human, ecosystem and animal health and well-being. Geneva, Switzerland: World Health Organization; 2021.
147. Martin AA. Why can't we control our food intake? The downside of dietary variety on learned satiety responses. *Physiol Behav*. 2016; 162: 120-129.