

Review

## Cytomegalovirus in Lung Transplant

Justin P. Rosenheck <sup>\*</sup>, Mena M Botros, David R Nunley

The Ohio State University Wexner Medical Center, 410 W. 10th Ave., Columbus, Ohio 43210, USA;  
E-Mails: [Justin.Rosenheck@osumc.edu](mailto:Justin.Rosenheck@osumc.edu); [Mena.Botros@osumc.edu](mailto:Mena.Botros@osumc.edu); [David.Nunley@osumc.edu](mailto:David.Nunley@osumc.edu)

\* **Correspondence:** Justin P Rosenheck; E-Mail: [Justin.Rosenheck@osumc.edu](mailto:Justin.Rosenheck@osumc.edu)

**Academic Editor:** Maricar Malinis

**Special Issue:** [Infections in Solid Organ Transplantation: 2020-2021](#)

*OBM Transplantation*

2021, volume 5, issue 2

doi:10.21926/obm.transplant.2102145

**Received:** March 15, 2021

**Accepted:** June 16, 2021

**Published:** June 27, 2021

### Abstract

Lung transplantation is a therapeutic option for patients with advanced lung diseases. Lung transplant outcomes have improved over time with improvements in the management of these complex patients. Cytomegalovirus is a common opportunistic organism affecting all solid organ transplant recipients. Characteristics unique to lung transplantation can make this virus difficult to manage, with myriad complications including graft failure and death. Ongoing research into and understanding of cytomegalovirus has opened exciting new avenues of management. We discuss the various manifestations of CMV related pathologies in the lung transplant recipient. We discuss current mainstays of risk stratification, diagnosis, and treatment, as well as present new and evolving concepts. Current medications are highly effective at preventing and treating CMV manifestations, but may be poorly tolerated. A new generation of therapies carry the promise of efficacy, with a greater safety profile and improved tolerance of adverse effects. We discuss host-virus immune interactions, specifically how these can be utilized in clinical practice to individualize the cytomegalovirus related care of lung transplant recipients. Finally, we turn our attention to the near horizon as we continue to evolve the care of this unique population.



© 2021 by the author. This is an open access article distributed under the conditions of the [Creative Commons by Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is correctly cited.

## **Keywords**

Cytomegalovirus; lung transplantation; review; immunity; treatment

## **1. Literature Review**

### **1.1 Introduction**

Cytomegalovirus (CMV) remains the most common opportunistic infection following lung transplantation, responsible for a substantial burden of morbidity and mortality [1]. Despite decades of increasing short and long-term lung transplantation success, CMV management has remained relatively static. However, new diagnostic, prophylactic and therapeutic strategies are becoming available and bring with them promise to improve the care of this immunosuppressed population. In this review, we discuss current standards of management for CMV related pathologies and examine how these standards may soon change.

### **1.2 Background**

CMV is a DNA virus, a member of the beta-herpesviridae family that is highly prevalent throughout populations. CMV is ubiquitous throughout society, with general seropositivity between 30 and 97%, and prevalence correlated with age and socioeconomic factors [2-4]. CMV may spread via close contact, body fluids exposure, sexually or perinatally [5, 6]. In immunocompetent individuals, initial (lytic phase) infection may be asymptomatic, or cause a febrile illness that can resemble Epstein-Barr Virus (EBV) mononucleosis [7]. Lytic phase infection leads to the synthesis of a number of CMV specific proteins, including immediate-early (IE), early, and late viral proteins [8], all of which have been exploited in developing new diagnostic tests to evaluate CMV specific immunity. The initial immune response to CMV involves innate immunity, before transitioning to an adaptive immune response characterized by humoral and CMV-specific T cell responses [9]. While early studies pointed to the prominence of the cytotoxic (CD8+) T-cell in controlling CMV infection, the CD4+ T-cell response is necessary for lasting viral control [10]. Regardless, after immune control, this virus, like other members of the Herpesviridae family, enters a state of latency. As latent virus tends to reside in monocytes, macrophages, lymphocytes, endothelial cells and bone marrow progenitor cells [9], viral DNA has consequently been isolated from myriad body fluids including saliva, urine, blood, and genital secretions [5]. In individuals who lose immune control of latent virus, CMV may reactivate, entering a lytic phase and potentially resulting in recurrent clinical disease.

As a group, lung transplant recipients are at an increased risk of CMV related disease compared to other solid organ transplant (SOT) recipients. Many factors exist to explain this finding. First, donor seroprevalence has increased over time by transplant era, increasing from 55.3% between January 1992 and December 2000 to 61.6% in the most recent era (January 2010 through June 2018). This increasing seroprevalence is perhaps due to increasingly aggressive donor acceptance practices, and an increase in mean donor age from 30 to 40 years over the same time period [11]. Secondly, when compared to kidney, liver, or heart, the lung may carry a higher burden of donor CMV infected

cells allowing more opportunity for reactivation [9]. Thirdly, compared to these other organ transplants, lung transplants tend to require the highest amounts of maintenance immunosuppression which may also increase the odds of recrudescence of viral disease. Finally, episodes of acute rejection in lung transplant recipients may necessitate high doses of corticosteroids, or lymphocyte depleting therapies which predispose recipients to increased risk of CMV reactivation [9].

Risk of CMV related complications can be stratified according to the serostatus of donor and recipient. Seronegative recipients receiving organs from seropositive donors are at greatest risk, followed by seropositive recipient of seropositive organs. In this instance, though seropositive recipients may have pre-transplant immunity to CMV, donors may still transmit novel CMV strains, or recipients may lose immunity to CMV over time [12]. Seropositive recipients of seronegative organs are at lesser risk, and finally, seronegative recipients of seronegative donor organs are at the lowest risk of developing CMV disease; their risk of de novo CMV viremia mirroring the general population. To this end, it may be important to follow CMV serostatus of this population to document their exposure and risk to CMV over time. These risk categories provide guidance for duration of post-transplant CMV prophylaxis. Recently, testing to determine the presence of recipient CMV immunity following lung transplant has been studied to better stratify risk of CMV reactivation and to personalize duration of CMV prophylactic measures [13]. While many of these studies have been performed in non-lung transplant populations, a few studies have looked specifically at lung transplant recipients and have yielded promising results.

The consequences of CMV infection are not minor. The most common direct end organ manifestations of CMV disease in lung transplant patients include CMV pneumonitis and CMV colitis. In each instance, management typically necessitates invasive diagnostics, histopathologic demonstration, and intravenous therapy with ganciclovir. CMV has also been shown to have indirect effects on the host, possibly by altering the host immune response with subsequent increases in rates of bacterial, fungal, and viral infections [14-16]. A correlation with chronic lung allograft dysfunction has been observed [17], though a direct causation has yet to be shown [18, 19]. CMV has been identified as the cause of death in about 1% of lung transplant recipients and a significant cause of graft failure during the first three years following transplant [1, 20-23]. In an early series of 59 lung transplant recipients not receiving CMV prophylaxis, 32 (54%) patients developed CMV infection, including 95% of CMV seropositive recipients [24]. Survival and significant complications were worse in CMV seronegative recipients who developed infection. Additionally, post-transplant CMV infection is an economic burden on health care systems in terms of cost and number of hospital days [25-27].

## **2. CMV Manifestations**

Cytomegalovirus (CMV) infection or viremia differs from CMV syndrome or disease. Viremia is defined as the detection of CMV in a tested specimen. While viremia in the most traditional sense may be defined as CMV isolated by culturing, this is rarely performed. Detection of antigens may be referred to as antigenemia. Most commonly, viral DNA is detected using PCR based techniques in blood or serum samples, and isolation of viral nucleic acid sequences is referred to DNAemia or RNAemia [28].

CMV syndrome is a definition exclusively used in SOT patients. This is defined as detection of CMV in the blood and at least 2 of the following criteria:

1. Fever  $\geq 38^{\circ}\text{C}$  for at least 2 days.
2. New or worsening fatigue.
3. Leukopenia or neutropenia on 2 separate measurements at least 24 hours apart.
4.  $\geq 5\%$  atypical lymphocytes.
5. Thrombocytopenia defined as a platelet count of  $<100,000$  cells/ $\mu\text{L}$ .
6. Transaminitis  $\geq 2$  times the upper limit of normal [28].


CMV may manifest differently in individual patients. Those who are immunocompetent are generally asymptomatic or may present with mononucleosis syndrome. While mononucleosis is commonly associated with EBV, there are some key similarities and differences between these viral syndromes. Both may present with protracted fevers, fatigue, and a maculopapular rash [29] with lymphocytosis and atypical lymphocytes. By contrast, CMV mononucleosis typically does not present with enlarged cervical lymph nodes or tonsillitis. It may cause exudative tonsillopharyngitis but is heterophile antibody negative [29, 30]. Immunocompromised patients, such as those with HIV, cancer, or SOT, may present with more severe manifestations.

CMV disease is broadly defined as viremia and end organ dysfunction resulting from viral infiltration. Ideally, CMV should be directly evidenced from affected tissue; typically, histopathologic changes are seen on biopsy material. CMV retinitis is an exception, as clinical diagnosis is sufficient without detection of CMV in a tissue sample. It is also recognized that high viral DNA levels in a tissue sample with negative serum levels may represent disease, although organ specific viral level cutoffs have not been established. The most common manifestations of CMV disease in lung transplant patients include CMV pneumonitis and CMV colitis. Additional diseases caused by CMV in both immunocompetent and immunocompromised individuals are also described.

## **2.1 Pulmonary**

Clinical features of CMV pneumonitis include fever, fatigue, hypoxemia, dyspnea at rest or exertion, tachypnea with a cough that is typically non-productive. Pulmonary infiltrates on imaging can accompany clinical findings and can include non-specific CT findings such as diffuse ground-glass opacities, small pulmonary nodules often in a military pattern, confluent consolidation, and interstitial reticulation without air space opacification. While not easily identified, the finding of large cells with the typical 'owl eye' intranuclear inclusion, also known as Cowdry inclusions, in tissue obtained at biopsy provides a conclusive diagnosis (Figure 1) [31]. And while viral isolation from bronchoalveolar lavage (BAL) is suggestive of infection, strict levels for diagnosis have not been established. BAL testing for CMV can be completed using culture and nucleic acid amplification and quantification. While low levels of CMV DNA in BAL fluid may represent asymptomatic viral replication, higher levels may indicate active disease. Various levels with varying sensitivity and specificity have been reported [32-37], and though there is no specific threshold, it is generally accepted that viral copies  $> 500,000/\text{mL}$  are more likely associated with CMV pneumonitis [38]. Concurrent histopathology increases sensitivity and specificity of detection of CMV pneumonitis [34]. The higher the viral load, the more likely the patient is to have CMV identified on tissue biopsy, and the more likely the patient is to have clinical features of CMV pneumonia or pneumonitis. CMV pneumonitis may be life threatening in patients who are severely immunosuppressed, causing

severe hypoxemia and progressive respiratory failure by causing transplant failure or rejection. Pneumonitis secondary to CMV is seen less frequently now given improved detection along with modern prophylactic and treatment regimens. Incidence ranges from 3-6% in some reviews, with mortality reported as high as 35% [39]. It is important to note that asymptomatic CMV replication is thought to contribute to the development of bronchiolitis obliterans syndrome (BOS) [16, 40].



**Figure 1** Cowdry Inclusions: Haematoxylin and eosin stain of bronchial epithelial cells with CMV inclusions in nuclei and cytoplasm. Infected cells have an “Owl’s eye” appearance.

## **2.2 Gastrointestinal**

Gastrointestinal infection in the form of colitis presents with fever, fatigue, abdominal pain and cramping, diarrhea that is at times bloody, with inflammatory colitis and ulcerations noted on endoscopy. Cowdry inclusions can be seen on pathology [28, 41, 42]. CMV may also cause gastritis [43], esophagitis [44], ileitis [45], obstruction [46], and protein-losing hypertrophic gastropathy [47]. Diagnosis is made with endoscopy and biopsy proven CMV isolation from appropriate mucosa.

## **2.3 Cardiovascular**

CMV has been implicated in both venous thrombosis and coronary atherosclerosis, although causal relationships have not been established. Infection is thought to promote smooth muscle proliferation and increase uptake of lipoproteins, leading to atherosclerosis [48, 49]. Pericarditis and myocarditis have also been reported, and can be confirmed by pericardial or myocardial biopsy and isolation, culture, immunohistochemistry, nucleic acid analysis, or identification of CMV histological features [28, 50].

## **2.4 Hepatic and Pancreatic**

Although hepatitis is a known complication of infection, asymptomatic mild transaminitis is often seen in patients infected with CMV. Disease is diagnosed by liver function test abnormalities and biopsy [28]. Granulomatous hepatitis secondary to CMV has been documented [51, 52], as have portal vein thrombosis [53, 54] and pancreatitis [28, 55].

## **2.5 Neurologic**

CMV is known to cause encephalitis [56], ventriculitis, and Guillain-Barré syndrome (GBS) [57, 58]. Other described neurological deficits associated with CMV include transverse myelitis [59], neuropathies, and cranial nerve palsies [60]. Diagnosis requires relevant clinical symptomatology and viral isolation from cerebrospinal fluid (CSF), culture from nervous tissue biopsy, or encephalitis on electroencephalography (EEG) [28].

## **2.6 Ocular**

CMV is known to cause anterior uveitis in immunocompetent patients when latent CMV is reactivated. While there are differences in the presentation of CMV related chronic anterior uveitis, it is a clinical diagnosis with anterior chamber inflammation, elevated intraocular pressure, and iris atrophy [61]. Retinitis is typically absent in anterior uveitis. Retinitis is diagnosed clinically, however, if the diagnosis is in question, CMV in the vitreous fluid can help support the diagnosis [28].

## **2.7 Renal**

Renal transplantations account for the majority of solid organ transplants performed annually. CMV nephritis is a significant issue in the renal transplant population. While CMV nephritis may theoretically affect lung transplant recipients, there are no published reports of CMV nephritis in this population.

## **3. CMV Monitoring**

There are multiple risk factors for the development of CMV disease in SOT and specifically in LT recipients, prevention is key given high rates of infection and reactivation. Late onset CMV occurs in patients who develop CMV related issues such as DNAemia or end organ complications after they have completed their course of primary prophylaxis post transplantation. This clinical phenomenon is most commonly seen in D+/R- patients between 3-6 months following discontinuation of prophylaxis. Late-onset disease also occurs with greater frequency in patients receiving high levels of immunosuppression, in those with acute rejection, and is seen more in LT compared to other SOTs [12]. Several studies have evaluated duration of prophylaxis [22, 23, 40, 62, 63], and have shown that CMV infection and disease remains frequent, increases mortality, and is decreased using extended prophylaxis [64, 65]. Prevention in the form of CMV prophylaxis and preemptive therapy are discussed below.

### **3.1 Universal Prophylaxis**

Given the high frequency of disease and CMV reactivation following LT, universal prophylaxis is the standard of care for at risk patients following lung transplant. This entails administration of antiviral medication to recipients of seropositive organs, or recipients who themselves are seropositive, for a predetermined length of time. Intravenous ganciclovir and the oral prodrug valganciclovir have been studied for universal prophylaxis and are the most commonly used agents today [12]. Universal prophylaxis effectively prevents disease, is relatively easy to implement, may prevent other opportunistic infections, and may improve survival by preventing rejection secondary

to infection [32]. Universal prophylaxis has been shown to prevent CMV disease in 58-80% of patients [16, 66]. Seronegative recipients who receive transplants from seronegative donors do not typically receive CMV prophylaxis, though may receive anti-viral prophylaxis directed at other pathogens.

### **3.2 Pre-emptive Therapy**

Pre-emptive therapy entails regular CMV monitoring and initiation of antivirals when there is evidence of CMV DNAemia in order to prevent CMV disease [12, 67]. While universal prophylaxis reduces rates of CMV disease as well as that of other herpes viruses, drug costs, potential side effects of antivirals, and development of resistance remain reasons to consider pre-emptive therapy [12].

Prophylaxis against CMV infection in the lung transplant recipient is a central tenet of post-operative care. While debate exists as to the primacy of universal or pre-emptive prophylaxis strategies in hematopoietic and other solid organ transplants, given the risk of developing active CMV following lung transplant universal CMV prophylaxis is provided for all lung transplant recipients with donor or recipient seropositivity for CMV. Duration of CMV prophylaxis is guided by risk of developing active CMV replication in the host. Seronegative recipients transplanted with seropositive donor organs represent the highest risk group and typically receive CMV prophylaxis for a period of at least 12 months [12, 65, 68]. Seropositive recipients will typically receive CMV prophylaxis for a period of at least 6 months [15, 30]. Longer durations of CMV prophylaxis, including indefinite have been implemented at some centers for the highest risk patients, though significant risks include the selection for and breakthrough of resistant CMV strains [69, 70], and medication associated myelotoxicity [71, 72]. After periods of universal prophylaxis, recipients will undergo periods of close pre-emptive monitoring. In the post-prophylaxis period, patients remain at risk for development of a primary viremia, replication of previously latent virus, or infection with a new strain of CMV. Close monitoring enables transplant teams to rapidly identify CMV replication and initiate a therapeutic response.


## **4. CMV Testing**

Serologic testing and viral cultures are no longer recommended methods of testing for the presence of CMV given the lack of specificity. Molecular testing has made monitoring CMV status and testing for disease much easier. Quantitative tests include antigen staining of leukocytes and PCR assays. Antigen testing is semiquantitative, where higher levels of antigenemia correlate with likelihood and severity of disease [73]. This test, however, is labor intensive, is less sensitive than PCR testing, and becomes less sensitive in neutropenic patients [74].

PCR assays are more sensitive quantitative tests for CMV, where higher levels correlate with disease [75]. Previously, each laboratory had their own standards and cutoffs - extraction method, amplification target, probe, non-standardized quantification of secondary standards, and amplicon size previously led to intra-lab variability [76]. These, however, have now been standardized through the 2010 WHO international standard for CMV QNAT [12]. Changes in CMV DNA levels should be normalized to log scale, and significant changes are recognized as 1 log increases or decreases in viral copies per mL of sample [12, 77]. Either plasma or whole blood can be used for PCR testing, although consistency in sampling should be considered, as CMV DNA is detected at higher levels in

whole blood as compared to plasma. Whole blood testing has the advantage of detecting latent and replicating virus and is a more sensitive test, while CMV plasma levels are more specific for active DNAemia [12, 74]. It is important to also note that changes in viral loads are useful to determine viral burden and control. Decreasing viral loads, whether in serum or other samples, may indicate improving disease control.

#### 4.1 CMV Agents (Figure 2) [78]



**Figure 2** Mechanism of action of antiviral agents against cytomegalovirus: The viral UL97 kinase phosphorylates viral and host proteins involved in viral DNA synthesis. This enzyme enables the initial phosphorylation of ganciclovir (GCV), which is then further phosphorylated by cellular kinases to the active form, GCV-triphosphate (GCV-TP). GCV-TP incorporates into viral DNA where it impedes DNA chain elongation. Maribavir (MBV) also interferes with UL97 by inhibiting nuclear egress of viral particles, and may also limit the initial phosphorylation. Cellular kinases are also responsible for conversion of cidofovir (CDV) into its active state, which incorporates into viral DNA, and leading to termination of chain elongation. The viral DNA polymerase coded by UL54 produces DNA multimers. The activity of this enzyme is directly inhibited by foscarnet (FOS). DNA multimers are then cleaved to monomers and prepared for packaging by the viral terminase complex. Letermovir (LMV) interferes with this activity by inhibiting the UL56 subunit of the viral terminase complex [78].

Ganciclovir and Valganciclovir have proven to be the workhorses of CMV directed therapies, applied as first line agents for both prophylaxis and treatment of CMV. Ganciclovir was first introduced in 1988, and in addition to its role in organ transplantation has found application in the treatment of congenital CMV and CMV in other immunocompromised states [7]. Ganciclovir is an acyclic nucleoside analogue that is administered in an inactive state and undergoes phosphorylation

by viral UL97 protein kinase in CMV infected cells to ganciclovir monophosphate. Subsequent phosphorylations are mediated by cellular kinases, generating the active form of this drug - ganciclovir triphosphate. Ganciclovir triphosphate inhibits the viral UL54 encoded DNA polymerase by competing with deoxyguanosine as a substrate for this enzyme. Incorporation of ganciclovir triphosphate during DNA elongation results in the premature termination of CMV DNA synthesis [7]. Ganciclovir is available in IV form, dosing and duration are dictated by indication – treatment or prophylaxis. An oral form of ganciclovir is no longer available in the United States. Ganciclovir is excreted by glomerular filtration and active tubular secretion, and dose adjustment is necessary in impaired renal function. In addition to the need for intravenous access, ganciclovir usage is limited by side effects, most notably bone marrow toxicity, manifested as leukopenia, neutropenia, anemia, and thrombocytopenia [79].

Valganciclovir is an oral prodrug of ganciclovir, that once absorbed is metabolized to ganciclovir by hepatic and intestinal esterases [9]. When appropriately dosed, serum concentrations of valganciclovir can be equivalent to IV ganciclovir and prevent CMV viremia [80, 81]. Valganciclovir has been shown to be an effective agent for CMV prophylaxis and treatment of mild or moderate CMV related diseases [82]. Despite the efficacy of valganciclovir, IV ganciclovir remains the agent of choice for patients with severe manifestations of CMV, due to its superior bioavailability [12].

Cytomegalovirus can be refractory or resistant to treatment with ganciclovir. When CMV is susceptible to ganciclovir, appropriately dosed regimens should lead to a significant decline, defined as a >1 log<sub>10</sub> decrease in CMV viral load within 2 weeks of initiation. Patients with an increase in viral load are described as having probable refractory (less than 1 log<sub>10</sub> increase) or refractory (greater than 1 log<sub>10</sub> increase) CMV [12]. Detection of refractory CMV should lead to testing for CMV resistance. CMV resistance is diagnosed upon detection of mutations with genotypic testing of the UL97 and UL54 genes and can be low-level or high-level. CMV resistance is seen in up to 4% of solid organ transplant recipients [83]. Risk factors for CMV resistance include over-immunosuppression, insufficient ganciclovir dosing [84], and CMV donor positive-recipient negative mismatch status [83, 85]. Patients with ganciclovir resistant CMV have increased mortality, number of hospital days and decreased renal function when compared to patients treated for ganciclovir sensitive CMV [83].

In some instances, low-level ganciclovir resistance may be overcome by increased dosing of ganciclovir. However, in the setting of high-level ganciclovir resistance or severe CMV disease, alternative agents should be deployed to treat CMV viremia and disease [12]. Guideline recommended secondary agents include foscarnet and cidofovir [12]. Both of these agents demonstrate a broad spectrum of activity against DNA viruses. Cidofovir bypasses the need for UL97 mediated phosphorylation, relying on cellular kinases present in high levels in CMV infected cells. Once activated, cidofovir competes with deoxycytidine triphosphate as a competitive substrate for viral DNA polymerase and leads to termination of viral DNA elongation. Resistance to cidofovir occurs in the setting of viral DNA polymerase mutations and occurs at a frequency similar to that of ganciclovir. Foscarnet inhibits the activity of the UL54 encoded viral DNA polymerase by competing at the pyrophosphate binding site, inhibiting the elongation of DNA chains, and inhibiting DNA synthesis [13]. Both agents require IV infusion, and suffer from similar adverse effects profiles, most notably a high incidence of renal injury and electrolyte derangement, albeit via different mechanisms [7, 9]. Brincidofovir was developed as an oral prodrug of cidofovir, however a randomized control trial comparing this medication to placebo for CMV prophylaxis in stem cell

transplant patients showed increased mortality and a higher burden of serious adverse effects [86]. Subsequently, brincidofovir is no longer being developed for management of CMV related purposes [13].

CMV immunoglobulin (CMV-Ig) preparations contain a standardized amount of purified human antibody to CMV, and may serve as an adjunct to prophylactic or treatment protocols [87]. CMV-Ig has been shown to delay the onset of CMV DNAemia and disease in high risk lung transplant recipients [88], and may reduce the risk of chronic lung allograft dysfunction [89]. Additionally, CMV-Ig has been shown to be an effective addition to antivirals as rescue therapy for life threatening CMV diseases [90]. However, the utility of CMV-Ig is debated as data supporting its use are mostly single-center, retrospective reports and a potentially diminishing role in the current era of improved CMV detection and treatment [91]. A Cochrane review found that there was no benefit to the use of CMV-Ig in prophylaxis of CMV related outcomes in solid organ transplant recipients [92].

Recently, new options for management of CMV which seek to resolve some of the most vexing issues associated with the traditional therapies have become clinically available. Letermovir is a non-nucleoside inhibitor of the UL56 encoded DNA terminase complex. It prohibits cleavage of CMV DNA multimers into monomers, thereby inhibiting packaging of viral DNA into the virion. Letermovir was approved in 2017 by the FDA for prophylaxis of CMV infections in hematopoietic stem cell transplant patients [93]. An ongoing randomized control trial is comparing the efficacy of letermovir versus valganciclovir to prevent CMV disease in adult kidney transplant recipients [94]. Though not yet indicated in lung transplant recipients, this medication has been used when resistance or side effects limit the use of traditional agents [95, 96]. Letermovir may be especially useful in the setting of CMV that is cross resistant to ganciclovir, foscarnet and cidofovir, though resistance to letermovir has already been demonstrated [97]. Letermovir is eliminated through hepatic uptake, has mild described adverse effects, does not cause myelosuppression nor does it impair renal function, though an interaction with calcineurin inhibitors may be expected [9]. Importantly, letermovir's spectrum of anti-viral activity is limited to CMV, thus additional anti-viral therapies may be indicated during letermovir use.

Maribavir is another new agent, with a potential role in treating CMV related diseases. Maribavir is a selective inhibitor of the UL97 encoded protein kinase, acting as a competitive inhibitor of ATP binding [98], and impedes nuclear egress of viral capsids [99]. Early trials failed to show efficacy of maribavir, though these used a low dosing regimen. With higher dosing, a phase 2 trial in both HSCT and solid organ transplant patients has demonstrated non-inferiority of maribavir to valganciclovir for CMV prevention [99]. Though UL97 mutations can cause cross resistance to ganciclovir and maribavir [78, 100], maribavir may have a role in treating CMV infections that are refractory or resistant to first and second line therapies [101]. Maribavir is well tolerated, without significant myelosuppression or nephrotoxicity, though complaints of GI distress and dysgeusia have been relatively common. Maribavir is specific for CMV and EBV, and additional antiviral coverage may be indicated in select patients while on maribavir.

An intact T cell response to CMV is integral to long term control of this virus. As such, in patients with recurrent or refractory CMV infections, reduction of immunosuppression may be considered to allow for a degree of immune reconstitution. This of course increases the risk of allograft rejection. Adaptive T-cell transfer of CMV specific T cells has been used successfully in the hematopoietic stem cell transplant population for control of CMV [13]. This technique has also been used in solid organ transplantation, including lung transplant recipients, with success in recent years [102-104]. In stem

cell transplant recipients, donor T cells may be used to induce viral immunity, however this is not effective in solid organ transplants. In solid organ recipients, current efforts center around stimulation of recipient peripheral blood mononuclear cells with purified CMV proteins, in vitro enhancement of an immune response and subsequent reinfusion of newly CMV immune cells to generate CMV immunity [105]. Subsequent changes in recipient T cell repertoire are predictive of successful adoptive T cell transfer [106]. Transfer of third party CMV-specific T cells is another paradigm currently being explored [107]. To date, Adoptive T cell therapies in solid organ transplant recipients have been shown to be generally safe and well tolerated, with minimal serious adverse effects. This technique may also allow for resumption of full dose immunosuppression, which may have previously been decreased to allow a host response to CMV to develop. Continued research in the use of adoptive transfer of CMV specific immunity in solid organ transplant recipients is ongoing [108].

Alternative approaches to managing CMV infection are in varying stages of clinical readiness. An immunosuppression regimen including mTOR inhibitors has been shown to reduce the incidence of CMV viremia in transplant recipients, although the mechanisms underlying this association are poorly understood [109-112]. Leflunomide impairs virion assembly and inhibits pyrimidine synthesis and has been used off label for resistant and refractory CMV [113]. A monoclonal AB, RG7667 is undergoing clinical trials and has shown promise as potential therapy to prevent CMV infection in solid organ transplant recipients [114].

An effective vaccine against CMV would prove to be a holy grail in the management of transplant recipients, pre-empting the risks associated with CMV diseases and medication related adverse effects. To this end, efforts to develop a vaccine have persisted since the 1970's and have been increased since the turn of the century [115]. An ideal vaccine would provide long lasting immunity by stimulating humoral and cellular arms of the adaptive immune system and lead to development of memory responses that protect against the entire range of CMV strains. Development has been slowed by incomplete understanding of the CMV virus, including its latent state, its unique immune evasion capabilities, an incomplete understanding of necessary immune elements for protection, and lack of good animal models [116]. A myriad of vaccines is currently in varying stages of development, including live, non-living, vectored, and novel vaccines [117], including an mRNA vaccine candidate [118]. There is clearly an important need for a CMV vaccine, both in the transplant community and for prevention of congenital CMV [119, 120]. However, to date, no vaccine is currently in Phase 3 trials.

## **5. CMV Immunity and Immunity Testing**

An evolving understanding of immunity against CMV has opened a new phase of personalized management in the care of lung transplant recipients. Interactions between CMV and the host immune system are complex and dynamic. Following initial viremia, a robust immune response is triggered, eliciting activity from innate and adaptive immunity, including elements of both the humoral and cell mediated arms. Innate immune cells, including natural killer (NK) cells, and proteins such as toll like receptors, characterize the initial response to CMV [121]. CD4, CD8 and antibody producing cells are elicited in the adaptive immune response [122]. Intact cellular immunity provides long lasting protection from active CMV viremia.

Prior to transplant, the presence of a serologic antibody response to CMV indicates prior exposure and therefore likely immune control of CMV replication. Following transplantation, however, the host immune system's ability to control CMV infection may be impaired secondary to induction and maintenance immunosuppression regimens. Immune monitoring of the response to CMV represents a new avenue to personalize management of CMV treatment and prophylaxis. A number of unique tests assessing the T cell response to CMV have been developed and are in varying stages of clinical readiness. Principles of testing vary, though all tests include the in vitro stimulation of immune cells with CMV specific proteins and subsequent assessment of the immune response. ELISpot nonspecifically tests both CD4 and CD8 responses, while QuantiFERON tests primarily monitor CD8 responses [40, 123, 124]. Flow cytometry-based T-cell immune profiles provide specific values for both CD4 and CD8 cellular responses [125, 126]. Ongoing efforts to refine these tests, to best understand appropriate threshold values and when to apply these tests are underway [62]. In short, demonstration of the presence or absence of host immunity to CMV holds great promise to assist the clinical team with decisions to prolong or discontinue CMV therapies.

## **6. Future of CMV in LTx and Conclusion**

Cytomegalovirus has been a vexing concern for lung transplant recipients and their healthcare providers since the inception of routine transplantation. CMV remains a complex and incompletely understood virus and is a burden on patients and health care systems, with direct and indirect effects impacting patient's quality and quantity of life. Major advances have allowed practitioners to better prevent and treat CMV in lung transplant recipients. Increasingly sensitive testing allows us to identify early viremia and risk stratify our patients. New therapies exploit our improved understanding of the virus, are less toxic and easier to administer. The future appears bright, as new therapies emerge, diagnostic strategies evolve, and promising vaccine options continue to be developed.

## **Author Contributions**

JR and MB are responsible for the conception and drafting of this work. DN is responsible for critical revision of this work. All authors approve of the final version of this work.

## **Funding**

This work was completed with no funding.

## **Competing Interests**

The authors have declared that no competing interests exist.

## **References**

1. Chambers DC, Cherikh WS, Harhay MO, Hayes D, Hsich E, Khush KK, et al. The international thoracic organ transplant registry of the international society for heart and lung transplantation: Thirty-sixth adult lung and heart-lung transplantation report-2019; focus theme: Donor and recipient size match. *J Heart Lung Transplant*. 2019; 38: 1042-1055.

2. Staras SA, Dollard SC, Radford KW, Flanders WD, Pass RF, Cannon MJ. Seroprevalence of cytomegalovirus infection in the United States, 1988-1994. *Clin Infect Dis.* 2006; 43: 1143-1151.
3. Bate SL, Dollard SC, Cannon MJ. Cytomegalovirus seroprevalence in the United States: The national health and nutrition examination surveys, 1988-2004. *Clin Infect Dis.* 2010; 50: 1439-1447.
4. Cannon MJ, Schmid DS, Hyde TB. Review of cytomegalovirus seroprevalence and demographic characteristics associated with infection. *Rev Med Virol.* 2010; 20: 202-213.
5. Hasing ME, Pang XL, Mabilangan C, Preiksaitis JK. Donor cytomegalovirus transmission patterns in solid organ transplant recipients with primary infection. *J Infect Dis.* 2020; 223: 827-837.
6. Haidar G, Boeckh M, Singh N. Cytomegalovirus infection in solid organ and hematopoietic cell transplantation: State of the evidence. *J Infect Dis.* 2020; 221: S23-S31.
7. Gugliesi F, Coscia A, Griffante G, Galitska G, Pasquero S, Albano C, et al. Where do we stand after decades of studying human cytomegalovirus? *Microorganisms.* 2020; 8: 685.
8. Gandhi MK, Khanna R. Human cytomegalovirus: Clinical aspects, immune regulation, and emerging treatments. *Lancet Infect Dis.* 2004; 4: 725-738.
9. Meesing A, Razonable RR. New developments in the management of cytomegalovirus infection after transplantation. *Drugs.* 2018; 78: 1085-1103.
10. Lim EY, Jackson SE, Wills MR. The CD4+ T cell response to human cytomegalovirus in healthy and immunocompromised people. *Front Cell Infect Microbiol.* 2020; 10: 202.
11. Chambers DC, Zuckermann A, Cherikh WS, Harhay MO, Hayes D, Hsich E, et al. The international thoracic organ transplant registry of the international society for heart and lung transplantation: 37th adult lung transplantation report - 2020; focus on deceased donor characteristics. *J Heart Lung Transplant.* 2020; 39: 1016-1027.
12. Kotton CN, Kumar D, Caliendo AM, Huprikar S, Chou S, Danziger-Isakov L, et al. The third international consensus guidelines on the management of cytomegalovirus in solid-organ transplantation. *Transplantation.* 2018; 102: 900-931.
13. Stern A, Papanicolaou GA. CMV prevention and treatment in transplantation: What's new in 2019. *Curr Infect Dis Rep.* 2019; 21: 1-11.
14. L'Huillier AG, Ferreira VH, Ku T, Bahinskaya I, Kumar D, Humar A. Improving our mechanistic understanding of the indirect effects of CMV infection in transplant recipients. *Am J Transplant.* 2019; 19: 2495-2504.
15. Husni RN, Gordon SM, Longworth DL, Arroliga A, Stillwell PC, Avery RK, et al. Cytomegalovirus infection is a risk factor for invasive aspergillosis in lung transplant recipients. *Clin Infect Dis.* 1998; 26: 753-755.
16. Snyderman DR, Limaye AP, Potena L, Zamora MR. Update and review: State-of-the-art management of cytomegalovirus infection and disease following thoracic organ transplantation. *Transplant Proc.* 2011; 43: S1-S17.
17. Paraskeva M, Bailey M, Levvey BJ, Griffiths AP, Kotsimbos TC, Williams TP, et al. Cytomegalovirus replication within the lung allograft is associated with bronchiolitis obliterans syndrome. *Am J Transplant.* 2011; 11: 2190-2196.
18. Roman A, Manito N, Campistol JM, Cuervas-Mons V, Almenar L, Arias M, et al. The impact of the prevention strategies on the indirect effects of CMV infection in solid organ transplant recipients. *Transplant Rev.* 2014; 28: 84-91.
19. Sharples LD, McNeil K, Stewart S, Wallwork J. Risk factors for bronchiolitis obliterans: A

- systematic review of recent publications. *J Heart Lung Transplant*. 2002; 21: 271-281.
20. Kerschner H, Jaksch P, Karigl G, Popow-Kraupp T, Klepetko W, Puchhammer-Stöckl E. Cytomegalovirus DNA load patterns developing after lung transplantation are significantly correlated with long-term patient survival. *Transplantation*. 2009; 87: 1720-1726.
  21. Opelz G, Döhler B, Ruhstroth A. Cytomegalovirus prophylaxis and graft outcome in solid organ transplantation: A collaborative transplant study report. *Am J Transplant*. 2004; 4: 928-936.
  22. Monforte V, Sintes H, López-Gallo C, Delgado M, Santos F, Zurbano F, et al. Risk factors, survival, and impact of prophylaxis length in cytomegalovirus-seropositive lung transplant recipients: A prospective, observational, multicenter study. *Transpl Infect Dis*. 2017; 19: e12694.
  23. Johansson I, Mårtensson G, Nyström U, Nasic S, Andersson R. Lower incidence of CMV infection and acute rejections with valganciclovir prophylaxis in lung transplant recipients. *BMC Infect Dis*. 2013; 13: 1-10.
  24. Duncan AJ, Dummer JS, Paradis IL, Dauber JH, Yousem SA, Zenati MA, et al. Cytomegalovirus infection and survival in lung transplant recipients. *J Heart Lung Transplant*. 1991; 10: 638-646.
  25. McCarthy JM, Karim MA, Krueger H, Keown PA. The cost impact of cytomegalovirus disease in renal transplant recipients. *Transplantation*. 1993; 55: 1277-1282.
  26. Das A. Cytomegalovirus infection in solid organ transplantation: Economic implications. *Pharmacoeconomics*. 2003; 21: 467-475.
  27. Kelly J, Hurley D, Raghu G. Comparison of the efficacy and cost effectiveness of pre-emptive therapy as directed by CMV antigenemia and prophylaxis with ganciclovir in lung transplant recipients. *J Heart Lung Transplant*. 2000; 19: 355-359.
  28. Ljungman P, Boeckh M, Hirsch HH, Josephson F, Lundgren J, Nichols G, et al. Definitions of cytomegalovirus infection and disease in transplant patients for use in clinical trials. *Clin Infect Dis*. 2017; 64: 87-91.
  29. Klemola E, Von Essen R, Henle G, Henle W. Infectious-mononucleosis-like disease with negative heterophil agglutination test. Clinical features in relation to Epstein-Barr virus and cytomegalovirus antibodies. *J Infect Dis*. 1970; 121: 608-614.
  30. Begovac J, Soldo I, Presecki V. Cytomegalovirus mononucleosis in children compared with the infection in adults and with Epstein-Barr virus mononucleosis. *J Infect*. 1988; 17: 121-125.
  31. Yazigi F, Kahwash B, Al Sufiani F, Conces M, Prasad V, Kahwash S. Histopathologic identification and pattern recognition of common viral infections in the general pathology practice: An illustrated review. *Ibnosina J Med Biomed Sci*. 2016; 8: 28-51.
  32. Beam E, Germer JJ, Lahr B, Yao JD, Limper AH, Binnicker MJ, et al. Cytomegalovirus (CMV) DNA quantification in bronchoalveolar lavage fluid of immunocompromised patients with CMV pneumonia. *Clin Transplant*. 2018; 32: e13149.
  33. Lodding IP, Schultz HH, Jensen JU, Kirkby N, Perch M, Andersen C, et al. Cytomegalovirus viral load in bronchoalveolar lavage to diagnose lung transplant associated CMV pneumonia. *Transplantation*. 2018; 102: 326-332.
  34. Chemaly RF, Yen-Lieberman B, Castilla EA, Reilly A, Arrigain S, Farver C, et al. Correlation between viral loads of cytomegalovirus in blood and bronchoalveolar lavage specimens from lung transplant recipients determined by histology and immunohistochemistry. *J Clin Microbiol*. 2004; 42: 2168-2172.
  35. Boeckh M, Stevens-Ayers T, Travi G, Huang ML, Cheng GS, Xie H, et al. Cytomegalovirus (CMV)

- DNA quantitation in bronchoalveolar lavage fluid from hematopoietic stem cell transplant recipients with CMV pneumonia. *J Infect Dis.* 2017; 215: 1514-1522.
36. Boivin G, Olson CA, Quirk MR, Kringstad B, Hertz MI, Jordan MC. Quantitation of cytomegalovirus DNA and characterization of viral gene expression in bronchoalveolar cells of infected patients with and without pneumonitis. *J Infect Dis.* 1996; 173: 1304-1312.
  37. Lee HY, Rhee CK, Choi JY, Lee JW, Lee DG. Diagnosis of cytomegalovirus pneumonia by quantitative polymerase chain reaction using bronchial washing fluid from patients with hematologic malignancies. *Oncotarget.* 2017; 8: 39736-39745.
  38. Chemaly RF, Yen-Lieberman B, Chapman J, Reilly A, Bekele BN, Gordon SM, et al. Clinical utility of cytomegalovirus viral load in bronchoalveolar lavage in lung transplant recipients. *Am J Transplant.* 2005; 5: 544-548.
  39. Santos CA, Brennan DC, Yusef RD, Olsen MA. Incidence, risk factors and outcomes of delayed-onset cytomegalovirus disease in a large retrospective cohort of lung transplant recipients. *Transplantation.* 2015; 99: 1658-1666.
  40. Westall GP, Cristiano Y, Levvey BJ, Whitford H, Paraskeva MA, Paul E, et al. A randomized study of quantiferon CMV-directed versus fixed-duration valganciclovir prophylaxis to reduce late CMV after lung transplantation. *Transplantation.* 2019; 103: 1005-1013.
  41. Razonable RR, Humar A. Cytomegalovirus in solid organ transplant recipients-guidelines of the American society of transplantation infectious diseases community of practice. *Clin Transplant.* 2019; 33: e13512.
  42. Beswick L, Ye B, van Langenberg DR. Toward an algorithm for the diagnosis and management of CMV in patients with colitis. *Inflamm Bowel Dis.* 2016; 22: 2966-2976.
  43. Xiong X, Liu F, Zhao WL, Ji XH, Chen WG, Zou H, et al. Cytomegalovirus infective gastritis in an immunocompetent host misdiagnosed as malignancy on upper gastrointestinal endoscopy: A case report and review of literature. *Hum Pathol.* 2019; 92: 107-112.
  44. Li LX, Chakinala RC. Cytomegalovirus esophagitis. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK556110/>.
  45. Taniwaki S, Kataoka M, Tanaka H, Mizuno Y, Hirose M. Multiple ulcers of the ileum due to Cytomegalovirus infection in a patient who showed no evidence of an immunocompromised state. *J Gastroenterol.* 1997; 32: 548-552.
  46. Dinesh BV, Selvaraju K, Kumar S, Thota S. Cytomegalovirus-induced colonic stricture presenting as acute intestinal obstruction in an immunocompetent adult. *BMJ Case Rep.* 2013; 2013: bcr2013200944.
  47. Tard C, Madhi F, Verlhac S, Hagège H, Epaud R, Jung C. Protein-losing gastropathy associated with cytomegalovirus in two sisters - case reports and review of the literature. *Arch Pediatr.* 2019; 26: 232-235.
  48. High KP. Atherosclerosis and infection due to *Chlamydia pneumoniae* or cytomegalovirus: Weighing the evidence. *Clin Infect Dis.* 1999; 28: 746-749.
  49. Mawhorter SD, Lauer MA. Is atherosclerosis an infectious disease? *Cleve Clin J Med.* 2001; 68: 449-458.
  50. Campbell PT, Li JS, Wall TC, O'Connor CM, Van Trigt P, Kenney RT, et al. Cytomegalovirus pericarditis: A case series and review of the literature. *Am J Med Sci.* 1995; 309: 229-234.
  51. Clarke J, Craig RM, Saffro R, Murphy P, Yokoo H. Cytomegalovirus granulomatous hepatitis. *Am J Med.* 1979; 66: 264-269.

52. de la Serna-Higuera C, González-García M, Milicua JM, Muñoz V. Cytomegalovirus granulomatous hepatitis in an immunocompetent patient. *Gastroenterol Hepatol.* 1999; 22: 230-231.
53. Puccia F, Lombardo V, Giannitrapani L, Licata A, Mazzola G, Soresi M. Case report: Acute portal vein thrombosis associated with acute cytomegalovirus infection in an immunocompetent adult. *J Ultrasound.* 2017; 20: 161-165.
54. Squizzato A, Ageno W, Cattaneo A, Brumana N. A case report and literature review of portal vein thrombosis associated with cytomegalovirus infection in immunocompetent patients. *Clin Infect Dis.* 2007; 44: e13-e16.
55. Chan A, Bazerbachi F, Hanson B, Alraies MC, Duran-Nelson A. Cytomegalovirus hepatitis and pancreatitis in the immunocompetent. *Ochsner J.* 2014; 14: 295-299.
56. Baghban A, Malinis M. Ganciclovir and foscarnet dual-therapy for cytomegalovirus encephalitis: A case report and review of the literature. *J Neurol Sci.* 2018; 388: 28-36.
57. Leung J, Sejvar JJ, Soares J, Lanzieri TM. Guillain-Barré syndrome and antecedent cytomegalovirus infection, USA 2009-2015. *Neurol Sci.* 2020; 41: 885-891.
58. Orlikowski D, Porcher R, Sivadon-Tardy V, Quincampoix JC, Raphaël JC, Durand MC, et al. Guillain-Barré syndrome following primary cytomegalovirus infection: A prospective cohort study. *Clin Infect Dis.* 2011; 52: 837-844.
59. Fux CA, Pfister S, Nohl F, Zimmerli S. Cytomegalovirus-associated acute transverse myelitis in immunocompetent adults. *Clin Microbiol Infect.* 2003; 9: 1187-1190.
60. Kolson DL, Gonzalez-Scarano F. HIV-associated neuropathies: Role of HIV-1, CMV, and other viruses. *J Peripher Nerv Syst.* 2001; 6: 2-7.
61. Chan NS, Chee SP, Caspers L, Bodaghi B. Clinical features of CMV-associated anterior uveitis. *Ocul Immunol Inflamm.* 2018; 26: 107-115.
62. Paez-Vega A, Cantisan S, Vaquero JM, Vidal E, Luque-Pineda A, Lobo-Acosta M, et al. Efficacy and safety of the combination of reduced duration prophylaxis followed by immuno-guided prophylaxis to prevent cytomegalovirus disease in lung transplant recipients (CYTOCOR STUDY): An open-label, randomised, non-inferiority clinical trial. *BMJ Open.* 2019; 9: e030648.
63. Herrera S, Khan B, Singer LG, Binnie M, Chaparro C, Chow CW, et al. Extending cytomegalovirus prophylaxis in high-risk (D+/R-) lung transplant recipients from 6 to 9 months reduces cytomegalovirus disease: A retrospective study. *Transpl Infect Dis.* 2020; 22: e13277.
64. Humar A, Limaye AP, Blumberg EA, Hauser IA, Vincenti F, Jardine AG, et al. Extended valganciclovir prophylaxis in D+/R- kidney transplant recipients is associated with long-term reduction in cytomegalovirus disease: Two-year results of the IMPACT study. *Transplantation.* 2010; 90: 1427-1431.
65. Palmer SM, Limaye AP, Banks M, Gallup D, Chapman J, Lawrence EC, et al. Extended valganciclovir prophylaxis to prevent cytomegalovirus after lung transplantation: A randomized, controlled trial. *Ann Intern Med.* 2010; 152: 761-769.
66. Eid AJ, Razonable RR. New developments in the management of cytomegalovirus infection after solid organ transplantation. *Drugs.* 2010; 70: 965-981.
67. Preiksaitis JK, Brennan DC, Fishman J, Allen U. Canadian society of transplantation consensus workshop on cytomegalovirus management in solid organ transplantation final report. *Am J Transplant.* 2005; 5: 218-227.
68. Finlen Copeland CA, Davis WA, Snyder LD, Banks M, Avery R, Davis RD, et al. Long-term efficacy

- and safety of 12 months of valganciclovir prophylaxis compared with 3 months after lung transplantation: A single-center, long-term follow-up analysis from a randomized, controlled cytomegalovirus prevention trial. *J Heart Lung Transplant*. 2011; 30: 990-996.
69. Drew WL, Paya CV, Emery V. Cytomegalovirus (CMV) resistance to antivirals. *Am J Transplant*. 2001; 1: 307-312.
  70. Kotton CN. CMV: Prevention, diagnosis and therapy. *Am J Transplant*. 2013; 13: 24-40.
  71. Beam E, Lesnick T, Kremers W, Kennedy CC, Razonable RR. Cytomegalovirus disease is associated with higher all-cause mortality after lung transplantation despite extended antiviral prophylaxis. *Clin Transplant*. 2016; 30: 270-278.
  72. Wiita AP, Roubinian N, Khan Y, Chin-Hong PV, Singer JP, Golden JA, et al. Cytomegalovirus disease and infection in lung transplant recipients in the setting of planned indefinite valganciclovir prophylaxis. *Transpl Infect Dis*. 2012; 14: 248-258.
  73. Mengoli C, Cusinato R, Biasolo MA, Cesaro S, Parolin C, Palù G. Assessment of CMV load in solid organ transplant recipients by pp65 antigenemia and real-time quantitative DNA PCR assay: Correlation with pp67 RNA detection. *J Med Virol*. 2004; 74: 78-84.
  74. Clark NM, Lynch JP, Sayah D, Belperio JA, Fishbein MC, Weigt SS. DNA viral infections complicating lung transplantation. *Semin Respir Crit Care Med*. 2013; 34: 380-404.
  75. Almaghrabi RS, Omrani AS, Memish ZA. Cytomegalovirus infection in lung transplant recipients. *Expert Rev Respir Med*. 2017; 11: 377-383.
  76. Hirsch HH, Lautenschlager I, Pinsky BA, Cardeñoso L, Aslam S, Cobb B, et al. An international multicenter performance analysis of cytomegalovirus load tests. *Clin Infect Dis*. 2013; 56: 367-373.
  77. Dioverti MV, Razonable RR. Clinical utility of cytomegalovirus viral load in solid organ transplant recipients. *Curr Opin Infect Dis*. 2015; 28: 317-322.
  78. Piret J, Boivin G. Clinical development of letermovir and maribavir: Overview of human cytomegalovirus drug resistance. *Antiviral Res*. 2019; 163: 91-105.
  79. Inc. SP. Ganciclovir Injection. Package Insert. Schaumburg: Inc. SP.; 2019.
  80. Kiser TH, Fish DN, Zamora MR. Evaluation of valganciclovir pharmacokinetics in lung transplant recipients. *J Heart Lung Transplant*. 2012; 31: 159-166.
  81. Wiltshire H, Paya CV, Pescovitz MD, Humar A, Dominguez E, Washburn K, et al. Pharmacodynamics of oral ganciclovir and valganciclovir in solid organ transplant recipients. *Transplantation*. 2005; 79: 1477-1483.
  82. Asberg A, Humar A, Rollag H, Jardine AG, Mouas H, Pescovitz MD, et al. Oral valganciclovir is noninferior to intravenous ganciclovir for the treatment of cytomegalovirus disease in solid organ transplant recipients. *Am J Transplant*. 2007; 7: 2106-2113.
  83. Fisher CE, Knudsen JL, Lease ED, Jerome KR, Rakita RM, Boeckh M, et al. Risk factors and outcomes of ganciclovir-resistant cytomegalovirus infection in solid organ transplant recipients. *Clin Infect Dis*. 2017; 65: 57-63.
  84. Stevens DR, Sawinski D, Blumberg E, Galanakis N, Bloom RD, Trofe-Clark J. Increased risk of breakthrough infection among cytomegalovirus donor-positive/recipient-negative kidney transplant recipients receiving lower-dose valganciclovir prophylaxis. *Transpl Infect Dis*. 2015; 17: 163-173.
  85. Heliövaara E, Husain S, Martinu T, Singer LG, Cypel M, Humar A, et al. Drug-resistant cytomegalovirus infection after lung transplantation: Incidence, characteristics, and clinical

- outcomes. *J Heart Lung Transplant*. 2019; 38: 1268-1274.
86. Marty FM, Winston DJ, Chemaly RF, Mullane KM, Shore TB, Papanicolaou GA, et al. A randomized, double-blind, placebo-controlled phase 3 trial of oral brincidofovir for cytomegalovirus prophylaxis in allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2019; 25: 369-381.
  87. Cytogam. Cytomegalovirus immune globulin intravenous (human) package insert. Bern, Switzerland: Saol Therapeutics, Inc; 2020. Available from: <https://www.fda.gov/media/77671/download>.
  88. Lopez Garcia-Gallo C, García Fadul C, Laporta R, Portero F, Millan I, Ussetti P. Cytomegalovirus immunoglobulin for prophylaxis and treatment of cytomegalovirus infection in the (Val)ganciclovir era: A single-center experience. *Ann Transplant*. 2015; 20: 661-666.
  89. Ruttman E, Geltner C, Bucher B, Ulmer H, Höfer D, Hangler HB, et al. Combined CMV prophylaxis improves outcome and reduces the risk for bronchiolitis obliterans syndrome (BOS) after lung transplantation. *Transplantation*. 2006; 81: 1415-1420.
  90. Santhanakrishnan K, Yonan N, Callan P, Karimi E, Al-Aloul M, Venkateswaran R. The use of CMVig rescue therapy in cardiothoracic transplantation: A single-center experience over 6 years (2011-2017). *Clin Transplant*. 2019; 33: e13655.
  91. Rea F, Potena L, Yonan N, Wagner F, Calabrese F. Cytomegalovirus hyper immunoglobulin for CMV prophylaxis in thoracic transplantation. *Transplantation*. 2016; 100: S19-S26.
  92. Hodson EM, Jones CA, Strippoli GF, Webster AC, Craig JC. Immunoglobulins, vaccines or interferon for preventing cytomegalovirus disease in solid organ transplant recipients. *Cochrane Database Syst Rev*. 2007: CD005129. DOI: 10.1002/14651858.CD005129.pub2.
  93. Marty FM, Ljungman P, Chemaly RF, Maertens J, Dadwal SS, Duarte RF, et al. Letermovir prophylaxis for cytomegalovirus in hematopoietic-cell transplantation. *N Engl J Med*. 2017; 377: 2433-2444.
  94. Letermovir versus valganciclovir to prevent human cytomegalovirus disease in kidney transplant recipients (MK-8228-002). *ClinicalTrials.gov*; 2021. Available from: <https://clinicaltrials.gov/ct2/show/results/NCT03443869>.
  95. Cherrier L, Nasar A, Goodlet KJ, Nailor MD, Tokman S, Chou S. Emergence of letermovir resistance in a lung transplant recipient with ganciclovir-resistant cytomegalovirus infection. *Am J Transplant*. 2018; 18: 3060-3064.
  96. Aryal S, Katugaha SB, Cochrane A, Brown AW, Nathan SD, Shlobin OA, et al. Single-center experience with use of letermovir for CMV prophylaxis or treatment in thoracic organ transplant recipients. *Transpl Infect Dis*. 2019; 21: e13166.
  97. Veit T, Munker D, Kauke T, Zoller M, Michel S, Ceelen F, et al. Letermovir for difficult to treat cytomegalovirus infection in lung transplant recipients. *Transplantation*. 2020; 104: 410-414.
  98. Imlay HN, Kaul DR. Letermovir and maribavir for the treatment and prevention of cytomegalovirus infection in solid organ and stem cell transplant recipients. *Clin Infect Dis*. 2020: ciaa1713. DOI: 10.1093/cid/ciaa1713.
  99. Maertens J, Cordonnier C, Jaksch P, Poiré X, Uknis M, Wu J, et al. Maribavir for preemptive treatment of cytomegalovirus reactivation. *N Engl J Med*. 2019; 381: 1136-1147.
  100. Chou SW, Wu JY, Song KN, Bo TE. Novel UL97 drug resistance mutations identified at baseline in a clinical trial of maribavir for resistant or refractory cytomegalovirus infection. *Antiviral Res*. 2019; 172: 104616.

101. Papanicolaou GA, Silveira FP, Langston AA, Pereira MR, Avery RK, Uknis M, et al. Maribavir for refractory or resistant cytomegalovirus infections in hematopoietic-cell or solid-organ transplant recipients: A randomized, dose-ranging, double-blind, phase 2 study. *Clin Infect Dis*. 2019; 68: 1255-1264.
102. Brestrich G, Zwinger S, Fischer A, Schmück M, Röhmhild A, Hammer MH, et al. Adoptive T-cell therapy of a lung transplanted patient with severe CMV disease and resistance to antiviral therapy. *Am J Transplant*. 2009; 9: 1679-1684.
103. Holmes-Liew CL, Holmes M, Beagley L, Hopkins P, Chambers D, Smith C, et al. Adoptive T-cell immunotherapy for ganciclovir-resistant CMV disease after lung transplantation. *Clin Transl Immunology*. 2015; 4: e35.
104. Pierucci P, Malouf M, Glanville AR, Beagley L, Smith C, Khanna R. Novel autologous T-cell therapy for drug-resistant cytomegalovirus disease after lung transplantation. *J Heart Lung Transplant*. 2016; 35: 685-687.
105. Smith C, Beagley L, Rehan S, Neller MA, Crooks P, Solomon M, et al. Autologous adoptive T-cell therapy for recurrent or drug-resistant cytomegalovirus complications in solid organ transplant recipients: A single-arm open-label phase I clinical trial. *Clin Infect Dis*. 2019; 68: 632-640.
106. Smith C, Corvino D, Beagley L, Rehan S, Neller MA, Crooks P, et al. T cell repertoire remodeling following post-transplant T cell therapy coincides with clinical response. *J Clin Invest*. 2019; 129: 5020-5032.
107. Neuenhahn M, Albrecht J, Odendahl M, Schlott F, Dössinger G, Schiemann M, et al. Transfer of minimally manipulated CMV-specific T cells from stem cell or third-party donors to treat CMV infection after allo-HSCT. *Leukemia*. 2017; 31: 2161-2171.
108. Donor CMV specific CTLs in treating CMV reactivation or infection in participants who have undergone stem cell transplant or solid organ transplant. *ClinicalTrials.gov*; 2021. Available from: <https://clinicaltrials.gov/ct2/show/NCT03665675>.
109. Strueber M, Warnecke G, Fuge J, Simon AR, Zhang R, Welte T, et al. Everolimus versus mycophenolate mofetil de novo after lung transplantation: A prospective, randomized, open-label trial. *Am J Transplant*. 2016; 16: 3171-3180.
110. Rittà M, Costa C, Solidoro P, Sidoti F, Libertucci D, Boffini M, et al. Everolimus-based immunosuppressive regimens in lung transplant recipients: Impact on CMV infection. *Antiviral Res*. 2015; 113: 19-26.
111. Ghassemieh B, Ahya VN, Baz MA, Valentine VG, Arcasoy SM, Love RB, et al. Decreased incidence of cytomegalovirus infection with sirolimus in a post hoc randomized, multicenter study in lung transplantation. *J Heart Lung Transplant*. 2013; 32: 701-706.
112. Pascual J, Royuela A, Fernández AM, Herrero I, Delgado JF, Solé A, et al. Role of mTOR inhibitors for the control of viral infection in solid organ transplant recipients. *Transpl Infect Dis*. 2016; 18: 819-831.
113. Silva JT, Pérez-González V, Lopez-Medrano F, Alonso-Moralejo R, Fernández-Ruiz M, San-Juan R, et al. Experience with leflunomide as treatment and as secondary prophylaxis for cytomegalovirus infection in lung transplant recipients: A case series and review of the literature. *Clin Transplant*. 2018; 32: e13176.
114. Deng R, Wang YH, Maia M, Burgess T, McBride JM, Liao XC, et al. Pharmacokinetics and exposure-response analysis of RG7667, a combination of two anticytomegalovirus monoclonal antibodies, in a phase 2a randomized trial to prevent cytomegalovirus infection in high-risk

- kidney transplant recipients. *Antimicrob Agents Chemother.* 2018; 62: e01108-e01117.
115. Stratton KR, Durch JS, Lawrence RS. *Vaccines for the 21st century: A tool for decisionmaking.* Washington, US: National Academies Press; 2000.
116. Kawaguchi Y, Kimura H, Mori Y. *Human Herpesviruses.* Singapore: Springer Singapore; 2018.
117. Cui XL, Snapper CM. Development of novel vaccines against human cytomegalovirus. *Hum Vaccin Immunother.* 2019; 15: 2673-2683.
118. Moderna completes enrollment of cytomegalovirus (CMV) vaccine (mRNA-1647) phase 2 study. Cambridge, Massachusetts: Moderna, Inc; 2020 [cited 2021 February 23]. Available from: <https://investors.modernatx.com/news-releases/news-release-details/moderna-completes-enrollment-cytomegalovirus-cmv-vaccine-mrna>.
119. Diamond DJ, La Rosa C, Chiuppesi F, Contreras H, Dadwal S, Wussow F, et al. A fifty-year odyssey: Prospects for a cytomegalovirus vaccine in transplant and congenital infection. *Expert Rev Vaccines.* 2018; 17: 889-911.
120. Plotkin SA, Wang D, Oualim A, Diamond DJ, Kotton CN, Mossman S, et al. The status of vaccine development against the human cytomegalovirus. *J Infect Dis.* 2020; 221: S113-S122.
121. Watkins RR, Lemonovich TL, Razonable RR. Immune response to CMV in solid organ transplant recipients: Current concepts and future directions. *Expert Rev Clin Immunol.* 2012; 8: 383-393.
122. Molina-Ortega A, Martín-Gandul C, Mena-Romo JD, Rodríguez-Hernández MJ, Suñer M, Bernal C, et al. Impact of pretransplant CMV-specific T-cell immune response in the control of CMV infection after solid organ transplantation: A prospective cohort study. *Clin Microbiol Infect.* 2019; 25: 753-758.
123. Thompson G, Boan P, Baumwol J, Chakera A, MacQuillan G, Swaminathan S, et al. Analysis of the QuantiFERON-CMV assay, CMV viraemia and antiviral treatment following solid organ transplantation in Western Australia. *Pathology.* 2018; 50: 554-561.
124. Kumar D, Mian M, Singer L, Humar A. An interventional study using cell-mediated immunity to personalize therapy for cytomegalovirus infection after transplantation. *Am J Transplant.* 2017; 17: 2468-2473.
125. Rogers R, Saharia K, Chandorkar A, Weiss ZF, Vieira K, Koo S, et al. Clinical experience with a novel assay measuring cytomegalovirus (CMV)-specific CD4+ and CD8+ T-cell immunity by flow cytometry and intracellular cytokine staining to predict clinically significant CMV events. *BMC Infect Dis.* 2020; 20: 1-11.
126. Sester U, Thijssen S, van Bentum K, Neumann F, Kubuschok B, Sester M, et al. Rapid identification of preformed alloreactive T cells for use in a clinical setting. *Transplantation.* 2004; 78: 607-614.



Enjoy *OBM Transplantation* by:

1. [Submitting a manuscript](#)
2. [Joining in volunteer reviewer bank](#)
3. [Joining Editorial Board](#)
4. [Guest editing a special issue](#)

For more details, please visit:

<http://www.lidsen.com/journals/transplantation>